

北西太平洋上の大気中ラドン濃度の時空間変動 Temporal and spatial variations of Radon-222 in the western North Pacific

坪井 一寛^{1*}; 松枝 秀和¹; 澤 庸介¹; 丹羽 洋介¹; 村山 昌平²

TSUBOI, Kazuhiro^{1*}; MATSUEDA, Hidekazu¹; SAWA, Yousuke¹; NIWA, Yosuke¹; MURAYAMA, Shohei²

¹ 気象研究所, ² 産業技術総合研究所

¹Meteorological Research Institute, ²National Institute of Advanced Industrial Science and Technology

A new compact radon measuring system has been developed for high-resolution observation of low-level radon-222 (Rn) for the remote sites, in collaboration with the MRI and AIST. The Rn measuring system was installed at 4 stations of Minamitorishima (MNM), Yonagunijima (YON), Chichijima (CCJ) and Ryori (RYO) operated by Japan Meteorological Agency (JMA) since 2007. The Rn measurements clearly show that distinct seasonal variations as well as frequent episodic events with Rn enhancement peaks on a synoptic scale are successfully captured at all 4 stations. Although the seasonal cycles depended on the stations, significant correlations between the Rn and other trace gases were found for the most of the synoptic-scale events, indicating a large impact of widespread pollutions from the East-Asian countries on the regional air quality over the western North Pacific.

キーワード: ラドン

Keywords: Radon