

地上と衛星の統合観測による周北極域の植生フェノロジーの検出の高精度化 Multidisciplinary in situ and satellite observations for accurate detection of phenology in sub- and Arctic ecosystems

永井 信^{1*}; 鄭 峻介²; 小林 秀樹¹; 伊川 浩樹⁴; 中井 太郎³; 金 龍元⁴; 鈴木 力英¹; 杉本 敦子²
NAGAI, Shin^{1*}; TEI, Shunsuke²; KOBAYASHI, Hideki¹; IKAWA, Hiroki⁴; NAKAI, Taro³; KIM, Yongwon⁴; SUZUKI, Rikie¹; SUGIMOTO, Atsuko²

¹ 海洋研究開発機構地球環境変動領域, ² 北海道大学大学院地球環境科学研究院, ³ 名古屋大学地球水循環研究センター, ⁴ アラスカ大学国際北極圏研究センター

¹Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology, ²Graduate School of Env.Science / Faculty of Env.Earth Science Hokkaido University, ³Hydrospheric Atmospheric Research Center, Nagoya University, ⁴International Arctic Research Center, University of Alaska Fairbanks

To accurately evaluate the spatio-temporal variability of ecosystem functions and service in sub- and Arctic regions under rapid meteorological and climate changes, global, long-term, and comprehensive phenological observations are required. Towards this aim, satellite remote-sensing is useful to detect the spatio-temporal variability of plant phenology such as the timing of start (SGS) and end of growing season (EGS). However, from the in situ ecological research viewpoint, the satellite remote-sensing has not been sufficiently tested and validated by ground-truthing. Here, (1) we performed daily field observations with time-lapse digital cameras in boreal forests in Alaska and Siberia; (2) we examined the relationship between satellite-observed vegetation indices and plant phenology; and (3) we evaluated the spatio-temporal variability of the timing of SGS and EGS in sub- and Arctic regions by using MODIS Terra and Aqua-observed green-red vegetation index (GRVI). We found that (1) satellite-observed vegetation indices (i.e. NDVI, EVI, and GRVI) mainly detected the plant phenology of forest floor in sparse forests; (2) large year-to-year variability of the timing of SGS was detected in eastern Siberia and western Ural Mountains, while that of EGS was not clearly detected; and (3) in contrast, large year-to-year variability of the timing of EGS was detected in western Alaska, which is mainly covered by tundra vegetation, while that of SGS was not clearly detected.

キーワード: フェノロジー, リモートセンシング, 陸域生態系, シベリア, アラスカ, 地上真値
Keywords: phenology, remote sensing, terrestrial ecosystem, Siberia, Alaska, ground-truthing