

針広混交林流域における水流出の年々変動 Changes in interannual variability of runoff in a conifer and deciduous hardwood mixed forested watershed

野口 正二^{1*}; 村上 亘¹; 谷 誠²
NOGUCHI, Shoji^{1*}; MURAKAMI, Wataru¹; TANI, Makoto²

¹ 森林総合研究所, ² 京都大学農学研究科

¹Forestry and Forest Product Research Institute, ²Graduate School of Agriculture, Kyoto University

The National Forest Management conducts forest management in National Forests for the fulfillment of multi functional roles of forest including long-term wood production management. On the other hand, there are few studies that evaluated the runoff characteristics including a state of the forest for a long term. This study was conducted within the Kamabuchi No1 experimental watershed (3.06ha) in North part of Japan. Hydrological observation has been continued in cold snowy region since 1939. It is the longest record in this region in Japan. The site is covered with Natural hardwood forest (ex. *Fagus crenata*, *Quercus mongolica* var. *grosseserrata* and *Quercus serrata*) and coniferous plantation forest (*Cryptomeria japonica* and *Chamaecyparis obtuse*) which planted around 1912 to 1916. Surficial geology is tuff and shaletic tuff of the Tertiary period and soils are clay loam. Meteorological observation was conducted Yamagata experimental forests located to 800m from the watershed to the northeast. A 71-year record (1939-2010) of the precipitation and runoff was used for an analysis of the flow-duration curve. Tree (DBH \geq 6cm) census in the watershed was also conducted at 5 times (1942, 1950, 1957, 1979, 2008). The tree volume of *Chamaecyparis obtuse* is a regular tendency and the tree volume of *Cryptomeria japonica* linearly increased. Stem volume of oak trees has increased remarkably from 1942 to 1979 but there was a close tendency of an increase in 2008 because mortality of oak trees occurred in the watershed. Based on 5 times tree census, positive linear relationship was found between tree volume and age of stand. While the proportion of plentiful runoff has shown a tendency to decrease over long term, those of ordinary, low and scanty runoffs have tended to increase with increasing the tree volume.

キーワード: 長期水文観測, 積雪寒冷地域, 流況曲線, 流出特性

Keywords: duration curve, cold snowy region, long term hydrological observation, runoff characteristics