

腕足動物における殻体タンパク質の網羅的同定 Comprehensive identification of shell matrix proteins in brachiopods

磯和 幸延^{1*}; 更科 功²; 大島 健志朗³; 紀藤 圭治⁴; 服部 正平³; 遠藤 一佳¹
ISOWA, Yukinobu^{1*}; SARASHINA, Isao²; OSHIMA, Kenshiro³; KITO, Keiji⁴; HATTORI, Masahira³; ENDO, Kazuyoshi¹

¹ 東京大学 理学系研究科, ² 東京大学 総合博物館, ³ 東京大学 新領域創成科学研究科, ⁴ 明治大学 農学部
¹Graduate School of Science, Univ. Tokyo, ²University Museum, Univ. Tokyo, ³Graduate School of Frontier Sciences, Univ. Tokyo, ⁴School of Agriculture, Meiji Univ.

Brachiopods are marine invertebrates that appeared in the Cambrian, and they have two shells like bivalves composed of calcium carbonate or calcium phosphate. Shells contain organic matrix, which have important roles in the biomineralization processes. Recently, many shell matrix proteins in molluscs have been identified, and their roles in shell formation have been discussed. On the other hand, shell matrix proteins in brachiopods have not been identified, except for partial amino acid sequences of a chromoprotein, named ICP-1. In this study, we performed comprehensive identification of shell matrix proteins of the brachiopod *Laqueus rubellus* using proteomics combined with transcriptomics. As a result, we identified a total of 18 shell matrix proteins. BlastP search showed that these proteins have no homologues in skeletal proteins identified from other phylum, suggesting that brachiopod and mollusc shells are different in origin.