

生痕化石によって乱された生痕化石：Chondrites と Phycosiphon に乱された Phymatoderma と，その古生態学的意義
Composite trace fossils: Phymatoderma reburrowed by Chondrites/Phycosiphon and its paleoecological implications

泉 賢太郎^{1*}
IZUMI, Kentaro^{1*}

¹ 東京大学大学院理学系研究科地球惑星科学専攻

¹Department of Earth and Planetary Science, University of Tokyo

Composite *Phymatoderma* specimens from the Pliocene deep-sea Shiramazu Formation in Japan, particularly those reburrowed by *Chondrites* and *Phycosiphon*, were analyzed to reveal the differences caused by the activities of these trace-makers. *Phymatoderma* reburrowed by *Phycosiphon* is significantly larger than non-reburrowed *Phymatoderma*, whereas *Phymatoderma* reburrowed by *Chondrites* shows no significant difference in burrow diameter compared with non-reburrowed *Phymatoderma*. The recognized size selectivity (i.e., preference for larger burrows) by the *Phycosiphon* trace-makers can be explained by considering the different feeding strategies of these two ichnogenera; namely deposit-feeding *Phycosiphon*-makers, which must have processed a significant mass of sediment to obtain sufficient organic matter, whereas chemosymbiotic *Chondrites*-producers, which did not require a lot of sediment to obtain nutrients. In order to test these interpretations, records of the Phanerozoic trace fossils reburrowed by *Chondrites/Phycosiphon* were compiled. Consequently, the *Phycosiphon* -preference toward relatively larger burrows was recognized, which supports the results of this study. The compilation also indicates that the burrow size has become a limiting factor for the *Phycosiphon*-producers that tried to rework the sediments within previous subsurface burrows, at least for 80 million years.