

A close relationship between global oceanic environmental changes and seafloor mineral deposition during the Phanerozoic

KATO, Yasuhiro^{1*}

¹Department of Systems Innovation, Graduate School of Engineering, University of Tokyo

Metal deposition on seafloor is strongly controlled by marine redox conditions. Fe-Mn and Mn oxide deposits are formed under oxygenated oceans. In striking contrast, Cu and Zn-bearing sulfide deposits are stable under anoxic oceans. Seafloor mineral deposits in turn are good indicators to redox conditions or redox changes of modern and ancient oceans.

There are numerous strata-bound ore deposits in the Japanese accretionary complexes. These deposits are mainly divided into three types; umber (Fe-Mn), Mn-rich, and volcanogenic massive sulfide (VMS; Besshi-type). The Mn-rich deposits are further divided into two subtypes that are associated with greenstone and NOT associated. Ages of these deposits provide us important constraints for a secular change of marine redox conditions over the past ~360 Myr. Depositional ages of umber and Mn deposits were previously determined by microfossils including radiolarians and conodonts. On the other hand, ages of the Besshi-type deposits are determined by Re-Os method (Nozaki et al., 2013). Oxide ore deposits such as umbers and Mn deposits were very likely precipitated in the modern-style oxygenated deep-sea. In contrast, Mn carbonate and VMS deposits were precipitated in the stagnant, O₂-deficient deep-sea during the Triassic and Jurassic periods. Seafloor mineral deposition closely related to global oceanic environmental changes may give us a hint for exploring the causes of mass extinction, and further for elucidating the evolution of life.

Nozaki, T., Y. Kato, K. Suzuki (2013) Late Jurassic ocean anoxic event: evidence from voluminous sulphide deposition and preservation in the Panthalassa. *Scientific Reports*, 3: 1889; doi:10.1038/srep01889.

Keywords: oceanic environmental change, seafloor mineral deposit, Japanese accretionary complexes, marine redox condition, Phanerozoic