

つくばにおける福島事故後6ヶ月間の大気中の放射性セシウム担体に関する研究 Study on the carrier of airborne radiocesium collected for six month in Tsukuba after the Fukushima nuclear accident

兼保直樹^{1*};小暮敏博²;向井広樹²;大橋英雄³;鈴木芙美恵³;赤田尚史⁴;奥田知明⁵;池盛文数⁶
KANEYASU, Naoki^{1*}; KOGURE, Toshihiro²; MUKAI, Hiroki²; OHASHI, Hideo³; SUZUKI, Fumie³; AKATA, Naofumi⁴
; OKUDA, Tomoaki⁵; IKEMORI, Fumikazu⁶

¹産業技術総合研究所, ²東京大学大学院理学研究科, ³東京海洋大学, ⁴核融合科学研究所, ⁵慶應義塾大学理工学部, ⁶名古屋環境科学調査センター

¹National Institute of Advanced Industrial Science and Technology, ²Graduate School of Science, The University of Tokyo, ³Tokyo University of Marine Science and Technology, ⁴National Institute for Fusion Science, ⁵Faculty of Science and Technology, Keio University, ⁶Nagoya City Institute for Environmental Sciences

To obtain the knowledge on the physico-chemical properties of airborne radionuclides, we had been collected size-resolved aerosol in Tsukuba, Japan, since April 28, 2011, although the data obtained do not include the first radioactive plumes that reached to Tsukuba on March 15, 2011. From the initial result, we proposed a hypothesis that the sulfate aerosol was the potential carrier of the ¹³⁴Cs and ¹³⁷Cs that had undergone the middle- to long-range transport from the damaged reactor. We further inferred that re-suspended soil particles that attached radionuclides were not the major airborne radioactive substances from late April to May, 2011 (Kaneyasu et al., 2012).

Nevertheless, there are some issues to be addressed on the nature of airborne radionuclides. Those are, a) until when the sulfate aerosol acted as a carrier of the radiocesium released from the reactor, or the other substances acted as carriers instead, and b) what is the carrier substance when the re-suspension or re-emission of became the dominant source in the airborne radiocesium.

In this study, we address these subjects by analyzing the long-term aerosol samples collected later than those presented in the previous study. The temporal change in the activity size distribution of radiocesium for six month will be discussed. In addition, the carrier substance of radiocesium in the coarse mode size range aerosol is investigated by use of the autoradiograph and scanning electron microscope to the aerosol sample collected in 2011 summer.

キーワード: 放射性セシウム, 粒径分布, 再飛散, 電子顕微鏡, オートラジオグラフ

Keywords: radiocesium, size distribution, re-suspension, electron microscope, autoradiograph