

細粒粘土鉱物に保持された微量セシウムの脱離挙動：セシウム固定に及ぼす粘土粒子凝集の影響

Desorption behavior of intrinsic cesium in smectite: Effect of aggregation on the cesium fixation in clay particles

福士 圭介^{1*}; 酒井 遥²; 糸野 妙子³; 田村 明弘³; 荒井 章司³

FUKUSHI, Keisuke^{1*}; SAKAI, Haruka²; ITONO, Taeko³; TAMURA, Akihiro³; ARAI, Shoji³

¹ 金沢大学環日本海域環境研究センター, ² 金沢大学理工学域自然システム学類, ³ 金沢大学大学院自然科学研究科自然システム学専攻

¹Institute of Nature and Environmental Technology, Kanazawa University, ²College of Science and Engineering, Kanazawa University, ³Graduate School of Natural Science and Technology, Kanazawa University

The radiocesium from the Fukushima Daiichi nuclear power plant accident is retained at the surface soils around the power plant. The expandable fine grained clay minerals such as smectite and vermiculate are the candidates for the host phases of radiocesium. The sorption mechanism of cesium in the clay minerals is expected to be cation exchange reaction in the interlayer of the clay minerals. Therefore, the retained Cs must be desorbed to the solutions in the presence of high concentrations of major cations. On the other hand, some natural observations after the Fukushima accident have shown that the radiocesium in the contaminated soils or sediments is merely desorbed to the water even in saline solutions (e.g. Aoi et al 2013 JPGU meeting). The purpose of the study is to reproduce the unexpected fixation of cesium in clay minerals from the laboratory experiment by using standard well characterized smectite (Kunipia-F). The desorption behavior of intrinsic trace Cs (10 nmol/g from LA-ICP-MS) in smectite by major cations were systematically examined. The results of the present study showed that the aggregation of smectite by the presence of the divalent cations or high concentration of monovalent cations lead to the fixation of cesium in the clay aggregates.

Keywords: cesium, smectite, desorption, fixation, aggregation