

バイカル湖ガスハイドレートに含まれる炭化水素ガスの水素同位体比 Hydrogen isotope of hydrate-bound hydrocarbons at Lake Baikal

八久保 晶弘^{1*}; 坂上 寛敏¹; 南 尚嗣¹; 山下 聡¹; 高橋 信夫¹; 庄子 仁¹; Khlystov Oleg²; Kalmychkov Gennadiy³; De Batist Marc⁴

HACHIKUBO, Akihiro^{1*}; SAKAGAMI, Hirotohi¹; MINAMI, Hirotsugu¹; YAMASHITA, Satoshi¹; TAKAHASHI, Nobuo¹; SHOJI, Hitoshi¹; KHLYSTOV, Oleg²; KALMYCHKOV, Gennadiy³; DE BATIST, Marc⁴

¹ 北見工業大学, ² ロシア科学アカデミー陸水学研究所, ³ ロシア科学アカデミー地球化学研究所, ⁴ ゲント大学

¹Kitami Institute of Technology, ²Limnological Institute, SB RAS, ³Vinogradov Institute of Geochemistry, SB RAS, ⁴Ghent University

Natural gas hydrates exist in sublacustrine sediments of Lake Baikal. Gas hydrates were first obtained from sub-bottom depths of 121 and 161 m in the Baikal Drilling Project well located at the southern Baikal basin. Recently, MHP (Multi-phase Gas Hydrate Project, 2009-2013) revealed distribution of gas hydrate in sub-bottom sediment at the southern and central Baikal basins. We obtained gas hydrate crystals from more than 25 places, and retrieved hydrate-bound gas onboard. We measured molecular and isotopic compositions of hydrate-bound gas.

According to the $\delta^{13}\text{C}$ - δD diagram for methane (Whiticar, 1999), high and low methane $\delta^{13}\text{C}$ values indicate thermogenic and microbial origins, respectively, and methane δD provides information on methyl-type fermentation or CO_2 reduction in the microbial field. Kida *et al.* (2006) and Hachikubo *et al.* (2010) reported that hydrate-bound methane of Lake Baikal was microbial origin via methyl-type fermentation, because methane δD was about -300 ‰. We found heavier methane ($\delta^{13}\text{C}$ ranged from -50 ‰ to -40 ‰) in the Kukuy Canyon area (central Baikal basin), indicating thermogenic origin. Methane δD was distributed from -330 ‰ to -270 ‰. Generally, δD of thermogenic methane of marine gas hydrates is much more heavier (more than -200 ‰). Methane δD of Lake Baikal gas hydrate seems to be about 100 ‰ smaller than that of marine gas hydrate. Matveeva *et al.* (2003) reported that δD of the lake bottom water was about -133 ‰. Possibly, methane δD of hydrate-bound methane derives from δD of water.

Hachikubo A, Khlystov O, Krylov A, Sakagami H, Minami H, Nunokawa Y, Yamashita S, Takahashi N, Shoji H, Nishio S, Kida M, Ebinuma T, Kalmychkov G, Poort J (2010) Molecular and isotopic characteristics of gas hydrate-bound hydrocarbons in southern and central Lake Baikal. *Geo-Mar Lett* **30**: 321-329. doi:10.1007/s00367-010-0203-1

Kida M, Khlystov O, Zemskaaya T, Takahashi N, Minami H, Sakagami H, Krylov A, Hachikubo A, Yamashita S, Shoji H, Poort J, Naudts L (2006) Coexistence of structure I and II gas hydrates in Lake Baikal suggesting gas sources from microbial and thermogenic origin. *Geophys Res Lett* **33**: L24603. doi:10.1029/2006GL028296

Matveeva TV, Mazurenko LL, Soloviev VA, Klerkx J, Kaulio VV, Prasolov EM (2003) Gas hydrate accumulation in the subsurface sediments of Lake Baikal (Eastern Siberia). In: Woodside JM, Garrison RE, Moore JC, Kvenvolden KA (eds) Proc 7th Int Conf Gas in Marine Sediments, 7-11 October 2002, Baku, Azerbaijan. *Geo-Mar Lett* **23(3/4)**: 289-299. doi:10.1007/s00367-003-0144-7.

Whiticar MJ (1999) Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane. *Chem Geol* **161**: 291-314. doi:10.1016/S0009-2541(99)00092-3

キーワード: ガスハイドレート, 結晶構造, バイカル湖, メタン, 安定同位体

Keywords: gas hydrate, crystallographic structure, Lake Baikal, methane, stable isotope