

ガス中蒸発法におけるナノ粒子生成過程の高速偏光“その場”観察 High-speed polarized in-situ observation in a nucleation process of nanoparticles produced by the gas evaporation method

木村 勇気^{1*}; 三原 新²; 大沼 隼志²; 石塚 紳之介¹; 村山 健太¹; 塚本 勝男¹
KIMURA, Yuki^{1*}; MIHARA, Arata²; ONUMA, Takashi²; ISHIZUKA, Shinnosuke¹; MURAYAMA, Kenta¹; TSUKAMOTO, Katsuo¹

¹ 東北大学大学院理学研究科地学専攻, ² 株式会社フォトロン
¹Tohoku University, ²Photron

The gas evaporation method has been investigated for more than half a century since the Kubo effect was reported (1962). There have been many studies on the produced nanoparticles mainly using a transmission electron microscope, which have elucidated the different physical properties of nanoparticles from those in bulk. On the other hand, there have been almost no reports on nucleation in smoke related to crystal growth. Recently, we achieved in-situ observation of the nucleation process in smoke using a double-wavelength Mach-Zehnder-type interferometer, which can determine the temperature and pressure at the nucleation simultaneously. A series of experiments clearly showed that smoke particles condense homogeneously only in a very high supersaturated environment [1-3]. In a preliminary experiment using tungsten trioxide, the smoke particles condensed with a degree of supersaturation as high as $\sim 10^6$. In this process, since evaporant is continuously supplied into the surrounding of the evaporation source, the flow of smoke after the nucleation and growth of nanoparticles has been simply considered as a consecutive process. The nucleation and growth of smoke particles should be a rapid process (ms order) due to high supersaturation, so the concentration of the evaporated vapor drastically decreases. However, the details of the formation process remain unknown.

In this study, we attempted to visualize the nucleation of nanoparticles and motion of smoke using a high-speed polarization image sensor (Photron Inc.) to clarify the details of the nucleation process of smoke particles. Since the sensor itself has pixels with micro-polarizers, a phase shift interferogram can be obtained in less than a millisecond because of the lack of mechanical movement free, and can therefore be applied to rapid phenomena such as nucleation in vapor phase. Here, we show the preliminary results of homogeneous nucleation of tungsten oxide from vapor phase.

[1] Y. Kimura, H. Miura, K. Tsukamoto, C. Li, T. Maki, Interferometric in-situ observation during nucleation and growth of WO_3 nanocrystals in vapor phase, *Journal of Crystal Growth*, 316 (2011) 196-200.

[2] Y. Kimura, K. Tsukamoto, Interferometric observation of temperature distributions in the smoke experiment, *J. Jpn. Soc. Microgravity* 28 (2011) S9-S12.

[3] Y. Kimura, K. K. Tanaka, H. Miura, K. Tsukamoto, Direct observation of the homogeneous nucleation of manganese in the vapor phase and determination of surface free energy and sticking coefficient, *Crystal Growth & Design*, 12 (2012) 3278-3284.

キーワード: 核生成, 偏光高速カメラ, その場観察

Keywords: Nucleation, High-speed polarized camera, in-situ observation