

## Akebono observations of EMIC waves in the slot region of the radiation belts Akebono observations of EMIC waves in the slot region of the radiation belts

坂口 歌織<sup>1\*</sup>; 笠原 禎也<sup>2</sup>; 小路 真史<sup>3</sup>; 大村 善治<sup>4</sup>; 三好 由純<sup>3</sup>; 長妻 努<sup>1</sup>; 熊本 篤志<sup>5</sup>; 松岡 彩子<sup>6</sup>  
SAKAGUCHI, Kaori<sup>1\*</sup>; KASAHARA, Yoshiya<sup>2</sup>; SHOJI, Masafumi<sup>3</sup>; OMURA, Yoshiharu<sup>4</sup>; MIYOSHI, Yoshizumi<sup>3</sup>; NA-  
GATSUMA, Tsutomu<sup>1</sup>; KUMAMOTO, Atsushi<sup>5</sup>; MATSUOKA, Ayako<sup>6</sup>

<sup>1</sup> 情報通信研究機構, <sup>2</sup> 金沢大学総合メディア基盤センター, <sup>3</sup> 名古屋大学太陽地球環境研究所, <sup>4</sup> 京都大学生存圏研究所, <sup>5</sup> 東北大学大学院理学研究科地球物理学専攻, <sup>6</sup> 宇宙航空研究開発機構 宇宙科学研究所

<sup>1</sup>National Institute of Information and Communications Technology, <sup>2</sup>Information Media Center, Kanazawa University, <sup>3</sup>Solar-Terrestrial Environment Laboratory, Nagoya University, <sup>4</sup>Research Institute for Sustainable Humanosphere, Kyoto University, <sup>5</sup>Department of Geophysics, Graduate School of Science, Tohoku University, <sup>6</sup>Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency

We present a unique observation of electromagnetic ion cyclotron (EMIC) waves in the deep inner magnetosphere at  $L = 2.5-5$  made by the Akebono satellite at altitudes of 3,300-8,700 km. The mode conversion, i.e., L mode (He<sup>+</sup> band)→R mode (He<sup>+</sup> band)→L mode (O<sup>+</sup> band) was clearly identified from the equator to high latitudes. In addition, we found rising tone structures, recently identified as EMIC triggered emissions, which could lead to bursty precipitation of relativistic electrons. First, we estimated the ion composition ratio (H<sup>+</sup>, He<sup>+</sup>, O<sup>+</sup>) = (83%, 16%, 1%) from polarization analysis. Second, we estimated minimum resonant electron energies with the observed EMIC waves and triggered emissions to be 1?10 MeV. The satellite trajectory during the wave observation was primarily through the slot region of electron radiation belts. The collocation implies possible contribution of EMIC waves to formation of the slot region of radiation belts after a magnetic storm.

Keywords: EMIC wave, slot region of the radiation belt, mode conversion, triggered emission, ion composition ratio, Akebono satellite