

## Plan of large SPE search by the $^{14}\text{C}$ content measurement in Japanese trees for the past 5000 years

MIYAKE, Fusa<sup>1\*</sup> ; MASUDA, Kimiaki<sup>1</sup> ; HAKOZAKI, Masataka<sup>2</sup> ; NAKAMURA, Toshio<sup>2</sup> ; KIMURA, Katsuhiko<sup>3</sup>

<sup>1</sup>Solar-Terrestrial Environment Laboratory, Nagoya University, <sup>2</sup>Center for Chronological Research, Nagoya University, <sup>3</sup>Faculty of Symbiotic Systems Science, Fukushima University

Radiocarbon ( $^{14}\text{C}$ ) is produced by incoming cosmic rays to the Earth. Produced  $^{14}\text{C}$  becomes  $^{14}\text{CO}_2$  and is absorbed by trees by photosynthesis. Then, tree-rings record the past cosmic ray intensity. Rapid yearly increases in the  $^{14}\text{C}$  content have been detected for the period from AD 774 to AD 775 and from AD 993 to AD 994. Although some candidates for the cause of these cosmic-ray events have been considered, it has been considered that the solar activity (large SPE) is the most plausible cause.

There is the possibility that a lot of  $^{14}\text{C}$  increase events like the AD 775 one are hidden in the periods when the  $^{14}\text{C}$  content has not been measured with a 1-year resolution. If we detect such events, we are able to discuss a detailed occurrence rate of large SPE which is very important factor to prepare for future large SPEs.

We are planning to search for  $^{14}\text{C}$  increase events by the measurements of  $^{14}\text{C}$  content in Japanese trees for this 5000 years. In this thesis, we are going to explain the plan and problems.

Keywords: radiocarbon, tree-rings, cosmic-ray, SPE