

## Physical Picture of 2-1/2D Driven Collisionless Magnetic Reconnection Physical Picture of 2-1/2D Driven Collisionless Magnetic Reconnection

CHENG, Chio<sup>1\*</sup>; INOUE, S.<sup>2</sup>; ONO, Y.<sup>3</sup>; HORIUCHI, R.<sup>4</sup>  
CHENG, Chio<sup>1\*</sup>; INOUE, S.<sup>2</sup>; ONO, Y.<sup>3</sup>; HORIUCHI, R.<sup>4</sup>

<sup>1</sup>Plasma and Space Science Center, National Cheng Kung University, Taiwan, <sup>2</sup>Graduate School of Engineering, University of Tokyo, Japan, <sup>3</sup>Department of Advanced Energy, University of Tokyo, Japan, <sup>4</sup>National Institute of Fusion Science, Japan  
<sup>1</sup>Plasma and Space Science Center, National Cheng Kung University, Taiwan, <sup>2</sup>Graduate School of Engineering, University of Tokyo, Japan, <sup>3</sup>Department of Advanced Energy, University of Tokyo, Japan, <sup>4</sup>National Institute of Fusion Science, Japan

The physical picture of how electrons and ions flow, how the electric and magnetic fields change, and how particles gain energy will be presented for the 2-1/2D collisionless driven magnetic reconnection. The 2-1/2 dimensional collisionless reconnection studies are performed using the particle simulation PASMO code [1] and theoretical analysis. In particular, we will provide the physical mechanism of how the poloidal current (including the Hall current in the downstream region) is generated and how the electrostatic potential is produced in the poloidal plane. The physical picture of how the quadrupole magnetic field and electrostatic potential are generated in the 2-dimensional (poloidal) plane is different from the one presented by Uzdensky and Kulsrud.[2]

[1] H. Ohtani and R. Horiuchi, Plasma Fusion Res., 4, 024 (2009)

[2] D. A. Uzdensky and R. M. Kulsrud, Phys. Plasma, 13, 062305 (2006)

キーワード: magnetic reconnection, numerical simulation, space plasma, laboratory experiment  
Keywords: magnetic reconnection, numerical simulation, space plasma, laboratory experiment