

立方晶カルシウムペロブスカイトの小さな剛性率 Small shear modulus of cubic CaSiO₃ perovskite

河合 研志^{1*}; 土屋 卓久²

KAWAI, Kenji^{1*}; TSUCHIYA, Taku²

¹ 東京工業大学大学院理工学研究科地球惑星科学専攻, ² 愛媛大学 GRC

¹Department of Earth and Planetary Sciences, Tokyo Institute of Technology, ²Geodynamics Research Center, Ehime University

Ca-perovskite (CaPv) is considered to be one of the most abundant minerals in the Earth's lower mantle (LM) and was suggested to have distinctly larger shear modulus than MgPv from static calculations and mean-field theory (Karki and Stixrude 1999; Stixrude et al. 2007). In this study the elasticity of cubic CaPv is reinvestigated using density functional constant-temperature first principles molecular dynamics simulations with strict calculation conditions. First, we computed the stable structure of CaPv and found that the cubic phase is more stable than the tetragonal and orthorhombic in the LM P,T condition. The thermal equation of state of CaPv was analyzed using the MD data set, which indicates its thermal properties including Gruneisen parameter quite similar to those of MgPv. Along the adiabatic temperature, CaPv was found to have higher density than the PREM and 12.5% iron-bearing MgPv. Next, we calculated elastic constants of cubic CaPv. Our new results clearly demonstrate that cubic CaPv does not have anomalously large shear modulus suggested by previous calculations with a small computation cell. This is because the cell applied in the previous studies is too small to allow the rotational phonon motion of SiO₆ octahedra related to the zone boundary optic phonon instability. Acoustic wave velocities were finally determined from the elastic moduli, indicating no significant differences in velocities between CaPv and iron-bearing MgPv.

キーワード: Ca ペロブスカイト, 弾性率, 下部マントル, 第一原理計算

Keywords: Ca-perovskite, elasticity, lower mantle, first principles