

長期海底地震観測で得られた2011年東北地方太平洋沖地震後の茨城・房総沖の地震活動

Spatial distribution of earthquakes off the coast of Ibaraki and the Boso Peninsula after the 2011 Tohoku Earthquake

中東 和夫^{1*}; 町田 祐弥²; 真保 敬⁷; 山田 知朗²; 望月 公廣²; 塩原 肇²; 篠原 雅尚²; 村井 芳夫³; 日野 亮太⁴; 東 龍介⁴; 鈴木 健介⁸; 久保田 達矢⁴; 長谷川 和也⁴; 佐藤 利典⁵; 高田 裕能⁵; 植平 賢司⁷; 八木原 寛⁶

NAKAHIGASHI, Kazuo^{1*}; MACHIDA, Yuya²; SHINBO, Takashi⁷; YAMADA, Tomoaki²; MOCHIZUKI, Kimihiro²; SHIOBARA, Hajime²; SHINOHARA, Masanao²; MURAI, Yoshio³; HINO, Ryota⁴; AZUMA, Ryosuke⁴; SUZUKI, Kensuke⁸; KUBOTA, Tatsuya⁴; HASEGAWA, Kazuya⁴; SATO, Toshinori⁵; TAKATA, Hiroyoshi⁵; UEHIRA, Kenji⁷; YAKIWARA, Hiroshi⁶

¹ 神戸大学, ² 地震研究所, ³ 北海道大学, ⁴ 東北大学, ⁵ 千葉大学, ⁶ 鹿児島大学, ⁷ 防災科学研究所, ⁸ 海洋研究開発機構

¹Kobe Univ., ²Earthquake Research Inst., ³Hokkaido Univ., ⁴Tohoku Univ., ⁵Chiba Univ., ⁶Kagoshima Univ., ⁷National Research Institute for Earth Science and Disaster Prevention, ⁸JAMSTEC

The 2011 off the Pacific coast of Tohoku Earthquake occurred on March 11, 2011, off shore of the northeast Japan region. Many aftershocks occurred following the mainshock. To obtain a precise aftershock activity is important for understanding the mechanism of earthquake generation, and the recovery of plate coupling at a ruptured plate boundary. In order to study the aftershock activity, we had deployed 66 long-term ocean bottom seismometers (LTOBS) off the coast of Ibaraki and the Boso Peninsula from October 2011 to November 2012.

For hypocenter determination, we selected 1606 events whose epicenter catalog which the Japan Meteorological Agency for hypocenter determination. P- and S- wave arrival times were manually picked using the WIN system (Urave and Tsukada, 1991). Hypocenters were determined by the maximum-likelihood estimation technique (Hirata and Matsuura, 1987). The hypocenter location program used in this study is based one-dimensional structure with constant V_p/V_s ratio of 1.73. Because a sedimentary layer below the sea floor generally has a large V_p/V_s value, an adjustment of the station corrections is needed. To obtain the station correction, we used the following method. First, we located the hypocenter using the P- and S-wave arrival times with the assumed station correction values for the velocity structure used. The averaged differences between observed travel time and estimated travel times (O-C times) for each station were then calculated. The averaged O-C times were added to the previous station correction values, and the hypocenters were relocated. We repeated this procedure eleven times. After this procedure, the averaged O-C times were less than 0.1 s for both the P-wave and S-waves. We estimated 458 hypocenter locations with an error of less than 5 km in the horizontal direction and less than 3 km in depth by using LTOBS data.

Most of the hypocenter locations have a depth shallower than 40km. The earthquakes form a plane dipping landward in the study area. Comparing the hypocenter locations with crustal structures obtained by active seismic studies (e.g. Miura et al., 2003). Many events occurred along the plate boundary. We also compared the hypocenter locations with aftershock distribution of the seismic observation conducted immediately after 2011 Tohoku Earthquake (Shinohara et al., 2012). Shinohara et al., (2012) reported that the low seismicity region has seen at the shallow part of the plate interface in the off-Fukushima. On the other hand, our results showed the seismicity is not low at the same region. This difference may reflect the change of stress fields at a ruptured plate boundary.