

粒子法による複雑な自由表面を有する岩石の破壊進展解析  
Numerical study for failure behavior of rock masses including complex free-surfaces using a particle method

武川 順一<sup>1\*</sup>; 三ヶ田 均<sup>1</sup>; 後藤 忠徳<sup>1</sup>  
TAKEKAWA, Junichi<sup>1\*</sup>; MIKADA, Hitoshi<sup>1</sup>; GOTO, Tada-nori<sup>1</sup>

<sup>1</sup> 京都大学大学院工学研究科

<sup>1</sup> Kyoto Univ.

Failure of rock mass including complex free surfaces is of importance in many engineering and scientific fields. This paper applied an advanced discretization approach to simulate quasi-static failure of rock mass within a Hamiltonian particle method (HPM) framework. In HPM, a free surface is introduced in a simple way, just by removing or ignoring outer particles. This potential can be developed to discretize numerical models including complex free surfaces without the increment of time for pre-processing. In the present study, we developed the numerical simulator based on HPM with a staggered particle technique for simulating brittle failure and AE activities in rock mass with incorporating the elasto-plastic damage model. We, first, conducted uni-axial compressive tests for validating the effectiveness of our approach. Next, we adopted rectangular and circular disc specimens with a hole as complex free surface models. Our numerical results had good agreement with those from laboratory experiments. This suggests that HPM would be a method to simulate failure behavior of rock mass without time-consuming pre-processing.

キーワード: 粒子法, 破壊進展挙動, 岩石

Keywords: particle method, failure behavior, rock mass