

## Minor element partitioning and mineralogy in limpets from a CO<sub>2</sub> vent site Minor element partitioning and mineralogy in limpets from a CO<sub>2</sub> vent site

Nehrke Gernot<sup>1\*</sup>; LANGER Gerald<sup>2</sup>; SADEKOV Aleksey<sup>2</sup>; BAGGINI Cecilia<sup>3</sup>; RODOLFO-METALPA Riccardo<sup>4</sup>; HALL-SPENCER Jason<sup>3</sup>; Bijma Jelle<sup>1</sup>; ELDERFIELD Henry<sup>2</sup>  
NEHRKE, Gernot<sup>1\*</sup>; LANGER, Gerald<sup>2</sup>; SADEKOV, Aleksey<sup>2</sup>; BAGGINI, Cecilia<sup>3</sup>; RODOLFO-METALPA, Riccardo<sup>4</sup>; HALL-SPENCER, Jason<sup>3</sup>; BIJMA, Jelle<sup>1</sup>; ELDERFIELD, Henry<sup>2</sup>

<sup>1</sup>Biogeosciences, Alfred-Wegener-Institut, <sup>2</sup>University of Cambridge, <sup>3</sup>School of Marine Science and Engineering, University of Plymouth, <sup>4</sup>CoReUs, IRD, Centre IRD de Noumea

<sup>1</sup>Biogeosciences, Alfred-Wegener-Institut, <sup>2</sup>University of Cambridge, <sup>3</sup>School of Marine Science and Engineering, University of Plymouth, <sup>4</sup>CoReUs, IRD, Centre IRD de Noumea

Specimens of the patellogastropod limpet *Patella caerulea* were collected within and outside a CO<sub>2</sub> vent site at Ischia, Italy. The shells were sectioned transversally and scanned for polymorph distribution by means of confocal Raman microscopy. Minor element to calcium ratios were measured using laser-ablation-inductively-coupled-plasma-mass-spectroscopy (LA-ICPMS). Mg/Ca, Sr/Ca, and Li/Ca ratios were determined in calcitic as well as aragonitic parts of the shells. This approach allows for investigating the effects of the polymorph and the seawater carbonate chemistry on minor element partitioning separately.

キーワード: calcification, proxy, CO<sub>2</sub>, mineralogy, patellogastropod limpet  
Keywords: calcification, proxy, CO<sub>2</sub>, mineralogy, patellogastropod limpet