

白亜紀後期北米内陸海におけるスカフィテス科アンモノイド類の殻の破損に基づく捕食－被食関係の推定
Durophagous predation on scaphitid ammonoids in the Late Cretaceous Western Interior Seaway of North America

竹田 裕介^{1*}; 棚部 一成²; 佐々木 猛智²; Landman Neil H.³
TAKEDA, Yusuke^{1*}; TANABE, Kazushige²; SASAKI, Takenori²; LANDMAN, Neil H.³

¹ 東京大学大学院理学系研究科地球惑星科学専攻, ² 東京大学総合研究博物館, ³ アメリカ自然史博物館
¹Department Earth and Planetary Science, The University of Tokyo, ²The University Museum, the University of Tokyo, ³American Museum of Natural History

The study of the evolution of predator-prey interactions has contributed much to our understanding of the ecological background of biodiversity change through geological time, because they represent a driving force of natural selection. This study is the first to report a trend of predation intensity on scaphitid ammonoids from the Turonian to the Maastrichtian (Late Cretaceous) on the basis of analysis of ventral shell breakage in large samples from the U.S. Western Interior Province. Analysis of 835 adult specimens revealed ventral shell breakage in 50 specimens. In most of the damaged specimens, the breakage occurred in a preferred position at the rear part of the body chamber. Ventral breakage is rare in the Turonian specimens, whereas it is common in the Campanian and Maastrichtian specimens. The shell diameter of adult scaphitid ammonoids tends to increase with time. The position of the breakage and the absence of repairs indicate that the ventral breakage resulted from lethal predation. Based on the incidence of breakage and the size and shape of the breaks, possible predators include fish, reptiles, and cephalopods such as *Placenticeras*, *Eutrephoceras*, and coleoids. Our statistical analysis of ventral shell breakage indicates that the incidence of lethal predation increased in conjunction with an increase in adult shell size, suggesting that the body size of the prey was an important factor in predator-prey interactions. In addition, the predatory damage is more extensive in larger adults.