

## サハリン島沖ガスハイドレート胚胎域の間隙水溶存ガス Dissolved gas in pore water at the hydrate-bound field off Sakhalin Island

八久保 晶弘<sup>1\*</sup>; 竹谷 敏<sup>2</sup>; VERESHCHAGINA Olga<sup>3</sup>; 坂上 寛敏<sup>1</sup>; 南 尚嗣<sup>1</sup>; 山下 聡<sup>1</sup>; 高橋 信夫<sup>1</sup>; 庄子 仁<sup>1</sup>; JIN Young K.<sup>4</sup>; OBZHIROV Anatoly<sup>3</sup>  
HACHIKUBO, Akihiro<sup>1\*</sup>; TAKEYA, Satoshi<sup>2</sup>; VERESHCHAGINA, Olga<sup>3</sup>; SAKAGAMI, Hiroto<sup>1</sup>; MINAMI, Hirotsugu<sup>1</sup>; YAMASHITA, Satoshi<sup>1</sup>; TAKAHASHI, Nobuo<sup>1</sup>; SHOJI, Hitoshi<sup>1</sup>; JIN, Young K.<sup>4</sup>; OBZHIROV, Anatoly<sup>3</sup>

<sup>1</sup> 北見工業大学, <sup>2</sup> 産業技術総合研究所, <sup>3</sup> ロシア科学アカデミー太平洋海洋学研究所, <sup>4</sup> 韓国極地研究所  
<sup>1</sup>Kitami Institute of Technology, <sup>2</sup>National Institute of Advanced Industrial Science and Technology (AIST), <sup>3</sup>Pacific Oceanological Institute, FEB RAS, <sup>4</sup>Korea Polar Research Institute

We summarize characteristics of dissolved gas in pore water in sub-bottom sediments off Sakhalin Island, where gas hydrates were retrieved. Gas hydrates have been discovered at the northeastern, southeastern and southwestern Sakhalin Island in the cruises of LV31 (2003), LV32 (2003), LV36 (2005), and LV39 (2006) in the framework of hydro-Carbon Hydrate Accumulations in Okhotsk Sea (CHAOS) project, and those of LV47 (2009), LV50 (2010), LV56 (2011), LV59 (2012), LV62 (2013), and LV67 (2014) in the framework of Sakhalin Slope Gas Hydrate (SSGH) project. We retrieved sediment cores (184 cores in total) including gas hydrates (29 cores) in these cruises (R/V Akademik M. A. Lavrentyev).

The dissolved gases in the pore water were extracted according to a headspace gas method. Basically, 10 mL of sediment and 10 mL of saturated aqueous solution of NaCl with a small amount of benzalkonium chloride (preservative) were introduced into a 25 mL vial to create a 5 mL headspace. The headspaces were flushed with nitrogen or helium as an inert gas. The vials were then shaken thoroughly and stored overturned. Molecular and stable isotope compositions of these samples were measured in our laboratory (Kitami Institute of Technology).

The depths of sulfate-methane interface (SMI) distributed mainly 1-3 mbsf, and those of hydrate-bound core were shallower (30 cm to 1 m). The dissolved gases in pore water below the SMI depth were primarily methane (more than approximately 99% of the total hydrate-bound gas), although ethane, propane, CO<sub>2</sub>, and hydrogen sulfide were also detected. The molecular and isotopic compositions of dissolved gas in the pore water indicated that the sediment cores including gas hydrates contain <sup>13</sup>C-rich methane and relatively high concentrations of <sup>13</sup>C-rich ethane compared to other gas-rich sediment cores. The gas-rich sediment cores without gas hydrates are characterized by high C<sub>1</sub>/(C<sub>2</sub>+C<sub>3</sub>) ratios and <sup>13</sup>C-depleted methane and ethane. We suggested the idea that small amount of thermogenic gas controlled occurrence of gas hydrate in the subsurface sediment off Sakhalin Island.

キーワード: ガスハイドレート, 間隙水, 安定同位体, オホーツク海, サハリン島  
Keywords: gas hydrate, pore water, stable isotope, Sea of Okhotsk, Sakhalin Island