

## 小惑星ベスタの熱進化の数値シミュレーション Thermal evolution simulation of asteroid Vesta

野上 竜彦<sup>1\*</sup>; 城野 信一<sup>1</sup>

NOGAMI, Tatsuhiko<sup>1\*</sup>; SHIRONO, Sin-iti<sup>1</sup>

<sup>1</sup> 名古屋大学 大学院環境学研究科, <sup>2</sup> 名古屋大学 大学院環境学研究科

<sup>1</sup>Division of earth and planetary science, Nagoya university, <sup>2</sup>Division of earth and planetary science, Nagoya university

Vesta has been regarded as the parent body of the HED meteorites. From the observation of DAWN spacecraft, the uppermost layer of Vesta is composed from howardite ranging from 50km to 80km (Jutzi et al. 2013). It is known that the ratio of the number of eucrites to diogenites is around two. Based on these facts, rapidly cooled magma layer on Vesta should be more than 10km in thickness.

In this work, I studied the evolution of internal thermal evolution of Vesta due to heating of decay of <sup>26</sup>Al. I calculate the temperature distribution by solving numerically heat conduction equation. According to Formisano et al.(2013), if Vesta completed its formation within 1.4Ma from the injections of <sup>26</sup>Al into the solar nebula, the degree of silicate melting of inside Vesta exceeds 50 vol%. But in that work, convection and melt migration were not taken into account. These two mechanisms contribute to cool Vesta. So it is expected that the formation of Vesta is completed more early if these are taken into account. By the way, it is known that it takes about a few hundreds year for Vesta-size planet to complete its formation in planetary formation standard model.

As a convection model, I adopted the simple model of Kaula (1979). It was assumed that generated melt migrates the surface instantaneously, and the migrating melt to the surface was accounted as the rapidly cooled magma. There are two parameters in this study, including a (the percentage of melt migration) and t<sub>0</sub> (formation time of Vesta), and perform simulation taking into account the convection and melt migration.

As a result, convection and melt migration substantially change the evolution of internal thermal structure, and total eruptive volume of melt considerably depends on a and t<sub>0</sub>. The magma volume increases as a increases. On the other hand, the magma volume decreases as t<sub>0</sub> increases.

When t<sub>0</sub>=0, corresponding to no decay of <sup>26</sup>Al at the beginning, and a>0.3, the erupting magma layer of 10km in thickness is formed. When a=1, that is all melt is erupted, the erupting magma layer of 10km is formed if t<sub>0</sub><0.9Ma. According to these results, Vesta should be completed its formation within 0.9Ma after CAI formation, and more than 30% of generated melt should migrate the surface. But all generated melts aren't necessarily erupted, and if a<1, Vesta has to be completed more early.

Therefore, it is suggested that the formation time of Vesta should be more early than the estimate by Formisano et al.(2013), and that planetary formation standard model might have to be reviewed.