

Lithology of Cambrian to Ordovician radiolarian chert in Kazakhstan with special reference to trace fossils

*Yoshitaka Kakuwa¹, Olga I Nikitina², Tatiana J Tolmacheva³

1.Gas Hydrate Laboratory, Organization for the Strategic Coordination of Research and Intellectual Properties, Meiji University, 2.Sarpaev Institute of Geological Sciences, Almaty, Kazakhstan, 3.A.P. Karpinskii Russian Geological Research Institute, St. Petersburg, Russia

Introduction

Trace fossils have been examined in the upper Cambrian to upper Ordovician radiolarian bedded cherts in Australia, Scotland and Newfoundland to clarify both the evolution of benthic animals and the change in environment of pelagic realm (Kakuwa and John, 2001; 2010; Kakuwa, 2014a; Kakuwa, 2014b). The result is summarized that Darriwilian is the age when benthic animals colonized widely on the pelagic deep ocean bottom. The other important and well-dated successions of radiolarian bedded chert (ribbon radiolarite) in Kazakhstan are studied to verify the result.

Examined bedded chert belongs to the Burubaital Formation that exposed in the southwest of Lake Balkhash, southern Kazakhstan. Age constraints of the examined two successions of bedded chert are given by Tolmacheva et al. (2001; 2004; unpublished data) who studied conodont fossils in detail.

Outline of the examined successions

Locality 89101

Upper Cambrian and Lower Ordovician chert is measured as thick as 36 meters. The succession is divided into three. The lower part consists of pale gray crystalline chert of upper Cambrian. The middle part consists of red chert of Tremadocian, and the upper part consists of pale gray, red and black cherts of Tremadocian to Floian. Radiolarians and sponge spicules are observed under the microscope. Thin laminations consistently develop in the cherts of this locality.

Locality 9706

Lower to Middle Ordovician bedded chert continuously exposed as thick as 70 meters. The lower half of the section consists of pale to dark gray, black and red cherts. The upper half of this succession, Dapping and Darriwilian in age, consists of red chert with some intercalations of white crystalline chert.

Characteristic knobby bedding planes that are considered as trace fossils by Tolmacheva et al. (2001) develop in the middle to upper part of the section. Several beds and laminations of sandstone and oligomitic conglomerate (breccia) occur in the same horizon. The grains of sandstone and conglomerate are composed of angular to subrounded chert, and are cemented by microcrystalline quartz. Sponge spicules are closely associated with the clastic rocks.

Trace fossils

No convincing trace fossils are found in the Upper Cambrian to Tremadocian chert of the locality 89101, while trace fossils are found in the Darriwilian red chert of the locality 9706. Trace fossils found are horizontal tunnel on bedding planes. They are as long as 10 cm and a few millimeters in diameter. Some tunnels are bifurcated and some are sinuous. Curious tubes are densely crowded in cherts of some limited horizons. They run oblique, parallel and vertical to the bedding plane. The tubes are 3 to 8 mm in diameter and as long as 2 cm in compacted length. The tubes are filled with chalcedonic quartz. Various relieves on the knobby bedding planes include probable trace fossils and diagenetic deformations.

Implications of trace fossils

Trace fossils are found in the radiolarian bedded chert of Darriwilian in Kazakhstan. This result is common to the other examined areas of Australia, Scotland and Newfoundland, but the trace

fossils found in Kazakhstan are different from those found in the aforementioned areas.

Acknowledgement

This study is financially supported by the Ministry of Education, Culture, Sports, Science and Technology of Japan (No.24540496).

References

- Kakuwa, Y., Web, J., 2001. SEPM Special Publication No. 88, 267-276.
Kakuwa, Y., Web, J., 2010. Australian Journal of Earth Sciences 57, 615-625.
Kakuwa, Y., 2014a. Annual Meeting of The Sedimentological Society of Japan (Tsukuba).
Kakuwa, Y., 2014b. Japanese Geosciences Union Meeting 2015 (Chiba).
Tolmacheva, T.J., Danelian, T., Popov, L.E., 2001. Geology 29, 755-758.
Tolmacheva, T.J., Holmer, L., Popov, L., Gogin, I., 2004. Geological Magazine 141, 699-715.

Keywords: trace fossils, Cambrian, Ordovician, radiolarian chert, Kazakhstan, evolution

The origin of plate tectonics , the origin of the deep sea floor , elucidation of the driving force , with Multi- Impact Hypothesis

*Akira Taneko¹

1.SEED SCIENCE Lab.


From the fact that A Wegener is in North and South America's and Africa coastal profile of the continent to match, and advocates the continental drift hypothesis, I was trying to prove the continuity of the paleontology and geology in the grounds. Now it has become plate tectonics in the empirical observations such as inversion and seabed tape recorder hypothesis of seafloor spreading hypothesis and mantle convection hypothesis and of the Mid-Atlantic Ridge discoveries and transform fault of discoveries and geomagnetism is almost dogma. However, the driving force of continental drift, which failed to show is Wegener remains still a mystery even in plate tectonics. Wegener had been pointed out, the origin of the formation origin and plate boundary of the deep sea floor, which accounts for 70% (-5km), efforts to explore the origin of plate tectonics had been forgotten. Arc-shaped archipelago and the basin of origin is also remains a mystery. The new paradigm can clarify this all unified manner has been desired.

It was mentioned in the "elucidation of the Earth and the moon of the missing link" from the earth physics and solar system planets science by the multi-impact hypothesis by abduction. According to it, in addition to the origin of this plate tectonics, the origin of the formation and deep sea Yosoko of the month, the origin of the origin or Mercury or Pluto of eccentricity or Jupiter large red rice of the core, meteorites and even more was the origin and differentiation of the asteroid belt also of origin is a proposal of a new paradigm that can be unified elucidated. Abduction is the idea that conclusion, according to some hypotheses about possible if you can explain the multiple of the current situation the correctness of the hypothesis is guaranteed. In the hypothesis that there is a physical meaning, If it is correct ideas, dramatic progress can be obtained.



CERRA passed since the birth of the solar system up to about 40 billion years ago and track deformed by Jupiter perturbations, when the tearing with tension of Jupiter and the sun, CERRA and the earth had been differentiated coagulation. In this hypothesis that multiple mantle debris collides almost the same orbit around briefly to Earth, it became a repeated species large extinction of the rationale, and stripped of mantle and the sea of 5km depth by isostasy. When Darwin raised happened to collision mantle defect by isostasy, plate the surrounding crust was peeling is recessed concave, the boundary cracks to form an arcuate archipelago. Arc-shaped archipelago and Tethys Java island, etc. at the time of the formation of the sea, which was centered on the Pacific Ocean, trench arc connected to the outside of the arc-shaped archipelago plate shows the dive under the arc-shaped archipelago. Plate boundary is due to the cracking of the time in which a plurality of cross-sectional mantle lobe collides with the Earth. In this hypothesis , because the Earth is rotating on its axis , when the mantle was missing by the collision became convex in isostasy , moment of inertia in the earth that became non-uniform (unbalanced) , the driving force for the moment of inertia to the minimum occurs .

Only one of the hypothesis, that can be described as evolution unified all of the mystery is the result of the abduction.

Keywords: The origin of plate tectonics , The origin of the deep sea floor, Elucidation of the driving force, Multi- Impact Hypothesis, Bimodal frequency of earth height, The moon formed by collision of CERRA mantle Ruptured debris lobe



Origin of Deep Ocean floor by MIH and Isostasy 5km depth



Minimization of the moment
of inertia ► Driving force

Multi-Impact Hypothes by mantle Impactor

Reconsideration of sedimentary place of the Triassic Muyinhe Formation in the Changning-Menglian belt of Southwest China

*Tsuyoshi Ito¹, Xin Qian², Qinglai Feng²

1.Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, 2.China University of Geosciences, Wuhan

Siliceous rocks of the Triassic Muyinhe Formation in the Changning-Menglian belt in southwestern Yunnan Province in Southwest China had been considered to be pelagic deposits. We observed them and analyzed their geochemistry, and reconsidered the sedimentary place.

The observation revealed that the siliceous rocks are characterized by inclusion of abundant radiolarian test (e.g., *Triassocampe* Dumitrica, Kozur, and Mostler, *Pseudostylosphaera* Kozur and Mostler, *Eptingium* Dumitrica, and *Paroertlispongus*) and the lack of rhythmical bedding. The geochemical results are as follows: the samples have high concentrations of SiO₂; most of the samples were plotted in the non-hydrothermal field on the Al-Fe-Mn diagram; most of the samples were plotted in the continental margin field on the Fe₂O₃/TiO₂-Al₂O₃/(Al₂O₃+Fe₂O₃) and (La/Ce)N-Al₂O₃/(Al₂O₃+Fe₂O₃) diagrams. In addition, the samples show a flat rare earth element pattern normalized to North America shale composite.

These observational and geochemical results strongly suggest that the siliceous rocks are unlikely to represent pelagic deposits, indicating that the extent of the pelagic ocean basins in the Paleotethys during the Triassic is probably less than previously believed. These non-pelagic deposits may represent the closure stage of the Paleotethys.

Keywords: Geochemistry, Triassic, siliceous rock, radiolaria, Paleotethys, Changning-Menglian belt

Micrometeorites from Triassic and Jurassic bedded cherts of the Mino and Chichibu belts, Southwest Japan.

*Mitsutaka Miura¹, Tetsuji Onoue¹

1.Department of Earth and Environmental Sciences, Faculty of Science, Kumamoto University.

Micrometeorites, which are submillimeter-sized extraterrestrial particles that survive atmospheric entry, originate from dust-producing objects such as comets and asteroids. Ancient micrometeorites found in sedimentary rocks are of key interest as a historical record of meteoroid populations in the solar system. We report the recovery of well-preserved micrometeorites, older than 240 Ma, in radiolarian chert from the Chichibu Belt on Ajiro Island (Middle Triassic) and Mino Belt on Inuyama area (Late Triassic to Early Jurassic). These study sections consist of rhythmic alternations of chert and shale beds. Samples were mechanically crushed and passed through 1.0 mm mesh sieve until less than 5 g of the fine fraction was collected. We also used a hydrofluoric acid dissolution method for determining micrometeorite content of chert samples. However, no micrometeorites have been recognized using this method. Magnetic components were separated using the method for liquid-suspended particles with a magnetic field strength of ~500 mT. Micrometeorites were handpicked from the magnetic components. The collection of micrometeorites comprised 72 cosmic spherules, which are particles that totally melted during atmospheric entry. Analysis of the accretion rate for cosmic spherules reveals high accretion rates of small spherules in the Middle Triassic (Anisian) and Early Jurassic (Hettangian). However, the possible link between the enhancements in the accretion rate of cosmic spherules and variations in the flux of extraterrestrial matter to the Earth requires further scrutiny.

Keywords: micrometeorite, Middle Triassic, Early Jurassic, Chichibu Belt, Mino Belt, bedded chert

Marine osmium isotope record across the Middle-Upper Norian transition in the Upper Triassic deep-sea deposits from the Mino Belt, Japan

*Honami Sato¹, Tatsuo Nozaki¹, Tetsuji Onoue², Jun-Ichi Kimura¹, Qing Chang¹, Yutaro Takaya³, Yasuhiro Kato⁴, Katsuhiko Suzuki¹

1.Japan Agency for Marine-Earth Science and Technology, 2.Kumamoto University, 3.Waseda University, 4.The University of Tokyo

Instead of a single mass extinction at the Triassic-Jurassic boundary, extinctions in major pelagic groups, such as radiolarians and conodonts, occurred stepwise during the last 15 Myr of the Triassic. Although a marked diversity decline on such pelagic fauna began in the end of Middle Norian, the cause of this extinction has been uncertain. In order to assess the Middle to the Late Norian environmental changes in the Panthalassa Ocean, we examined secular changes in the marine osmium isotope compositions ($^{187}\text{Os}/^{188}\text{Os}$) recorded in the Triassic bedded chert succession of the Mino Belt, central Japan. The osmium isotope compositions of seawater reflect contributions of the riverine ($^{187}\text{Os}/^{188}\text{Os} \approx 1.4$), hydrothermal, and extraterrestrial ($^{187}\text{Os}/^{188}\text{Os} \approx 0.12-0.13$) inputs of osmium to the global ocean. Given the distinctive $^{187}\text{Os}/^{188}\text{Os}$ compositions of these inputs and the relatively short residence time of Os in the ocean (several tens of thousand years), seawater $^{187}\text{Os}/^{188}\text{Os}$ compositions are highly sensitive to changes in these fluxes. The Os isotope compositions show a gradual increase in $^{187}\text{Os}/^{188}\text{Os}$ during the Middle Norian. Two negative Os isotope excursions are observed within a ~2 m stratigraphic interval located between the Middle and the Upper Norian. The onset of the first negative Os excursion is roughly located at the base of the *Epigondolella bidentata* conodont zone in the late Middle Norian. The timing of this Os isotope excursion coincides with both the abrupt increase in Os concentration and low Re/Os ratios, all of which suggest a significant input of extraterrestrial Os into the sediments. The Os isotope compositions exhibit the second negative Os isotope excursion from the initial Os isotope composition of ~0.7 to unradiogenic values of ~0.4 in the early Late Norian. However, it is difficult to specify the cause for the second negative excursion with the dataset acquired in this study. Additional research is needed to clarify the cause for this record in the upper Norian.

Keywords: osmium isotope, chert, Upper Triassic, Mino Belt

Origin of the Lower Triassic deep-sea chert gap

*Shun Muto¹, Satoshi Takahashi¹, Satoshi Yamakita², Masayuki Ikeda³, Tetsuji Onoue⁴

1.Department of Earth and Planetary Science, Graduate School of Science, University of Tokyo,

2.Department of Earth Sciences, Faculty of Education and Culture, University of Miyazaki,

3.Department of Earth Science, Shizuoka University, 4.Department of Earth and Environmental Sciences, Kumamoto University

Carboniferous to Jurassic pelagic deep-sea sediments that accumulated in the pelagic Panthalassa Ocean are mostly composed of bedded chert lithofacies (Matsuda and Isozaki, 1991). Among them, the Lower Triassic interval is characterised by siliceous claystone dominant lithofacies instead of bedded chert lithofacies (the deep-sea chert gap) (Ishiga and Yamakita, 1993; Isozaki, 1997). The age of the deep-sea chert gap (ca. 252–247 Ma) corresponds to the time interval of ecosystem recovery from the end-Permian mass extinction, the largest mass extinction event in the Phanerozoic (Chen and Benton, 2012). Therefore, the deep-sea chert gap may reflect anomalous environmental conditions during the Early Triassic that possibly delayed the re-establishment of complex ecosystems after the mass extinction. The deep-sea chert gap may either be a result of an increase in burial flux of fine clastic material, or a decline in burial flux of biogenic silica in the Early Triassic pelagic Panthalassa. However, previous studies have not calculated burial fluxes of fine clastic material and biogenic silica based on well-dated sedimentary sequences. In order to identify the mechanism that led to the formation of the deep-sea chert gap, we estimated the burial fluxes of clastic material and biogenic silica in deep-sea sedimentary sequences, which were newly described in this study.

Detailed geological mapping and reconstruction of lithostratigraphy was carried out at Ogama in the Kuzuu area, Tochigi Prefecture, and in the Tsukumi area, Oita Prefecture, and consequently, several continuous sections were identified. Based on conodont biostratigraphy, the studied sections are correlated to the *triangularis-collinsoni* Zone, the *homeri* Zone, the *timorensis* Zone and the *bulgarica* Zone of Koike (1981), which corresponds to the Spathian (upper Olenekian; uppermost Lower Triassic) to middle Anisian (lower Middle Triassic). Using the conodont biostratigraphic framework, absolute ages of volcanic tuff beds in South China were projected onto the reconstructed lithostratigraphic columns of the deep-sea sections. Consequently, the lower limit of the linear sedimentation rate (LSR) was obtained for the Spathian and lowermost Anisian siliceous claystone dominant lithofacies.

In addition, major element concentrations were measured for rock samples obtained from the studied sections by X-ray fluorescence (XRF) analysis. The SiO_2 and Al_2O_3 concentrations were combined with measured rock density data and the lithostratigraphic data of the studied sections to calculate the burial fluxes of clastic material and biogenic silica for the stratigraphic intervals for which LSR was estimated. For the lowermost Anisian in Ogama, the minimum burial fluxes of clastic material and biogenic silica are $1030 \text{ g/cm}^2\cdot\text{m.y.}$ and $1040 \text{ g/cm}^2\cdot\text{m.y.}$, respectively. For the Spathian in the Tsukumi area, the minimum burial fluxes of clastic material and biogenic silica are $960 \text{ g/cm}^2\cdot\text{m.y.}$ and $1130 \text{ g/cm}^2\cdot\text{m.y.}$, respectively. The estimated minimum burial flux of clastic material for the deep-sea chert gap greatly exceeds the burial flux calculated for the Anisian bedded chert lithofacies in the Tsukumi area ($80 \text{ g/cm}^2\cdot\text{m.y.}$), calculated based on LSR by Soda et al. (2015). The estimated minimum burial flux of biogenic silica for the deep-sea chert gap is also higher than that of Anisian bedded chert in the Tsukumi area ($390 \text{ g/cm}^2\cdot\text{m.y.}$), calculated based on LSR by Soda et al. (2015). These results indicate that, during the deposition of sediments in the deep-sea chert gap, burial fluxes of clastic material and biogenic silica were both elevated, but

the increase is greater in the former. Hence, the deep-sea chert gap is a result of anomalously high flux of clastic material to the pelagic realm, which diluted biogenic silica in deep-sea sediments.

Keywords: Lower Triassic, deep-sea chert gap, conodont, clastic burial flux, biogenic silica burial flux

Phylogeny of Middle Jurassic *Striatojaponocapsa* and middle Cretaceous *Turbocapsula*, and potential of reliable radiolarian zonation

*Atsushi Matsuoka¹, Xin Li²

1.Department of Geology, Faculty of Science, Niigata University, 2.Graduate School of Science and Technology, Niigata University

The four-segmented *Stichocapsa tegiminis* Yao group gave rise to three-segmented *Striatojaponocapsa plicarum* (Yao). The evolutionary first appearance biohorizon defines the base of the *Striatojaponocapsa plicarum* Zone of Bajocian-early Bathonian age. In turn, *Striatojaponocapsa plicarum* (Yao) evolved into *Striatojaponocapsa conexa* (Matsuoka) through *Striatojaponocapsa synconexa* O'Dogerty, Gorican and Dumitrica. The evolutionary first appearance biohorizon of *Striatojaponocapsa conexa* defines the base of *Striatojaponocapsa conexa* Zone of late Bathonian-early Callovian age. This is a well-known evolutionary lineage of Middle Jurassic radiolarians and contributes to offer a reliable chronological framework. An evolutionary trend similar to the *Striatojaponocapsa* lineage is recognized in middle Cretaceous radiolarians belonging to the genus *Turbocapsula*. Four-segmented forms of *Turbocapsula* evolved into three-segmented *Turbocapsula figitiva* O'Dogerty by reducing segment number. *Turbocapsula figitiva* gave rise to *Turbocapsula costata* (Wu) by acquiring widely-spaced costae. This speciation is characterized by cladogenesis. The *Turbocapsula* lineage has a potential to produce a reliable radiolarian zonal scheme for the middle Cretaceous (Barremian-Aptian).

Keywords: radiolaria, phylogeny, zonation

Lithostratigraphy and radiolarian age of pelagic sediments in the Wadi Hilti area of the Oman Ophiolite: age constraint for eruption of the V2 lava

*Toshiyuki Kurihara¹, Kousuke Hara¹, Yumi Agui¹

1. Graduate School of Science and Technology, Niigata University

The Oman Ophiolite is the most complete preserved section of upper mantle to oceanic crust worldwide (e.g., Lippard et al., 1986). The ophiolite forms a huge thrust sheet that crops out for a length of 500 km and a width of approximately 80 km in the Oman Mountains. Ernewein et al. (1988) summarized the subdivision of volcanic rocks and recognized that V1, which has a normal MORB (N-MORB)-like signature, was erupted around a spreading ridge and that V2 formed by intra-oceanic arc volcanism. In addition, thick lava flows of V3, ascribed to intra-plate volcanism, were emplaced onto pelagic sediments (Umino, 2012).

The V2 and V3 lavas are widely distributed in the Wadi Hilti area, about 25 km west of Sohar, northern Oman Mountains. Recently, Kusano et al. (2014) reexamined the volcano-stratigraphy of the V2 lava in this area, consisting of lower LV2 (IAT) and upper UV2 (boninite) units, based on their geochemical composition and stratigraphic relationship. The eruption and emplacement mechanism of the V3 lava were studied by Umino (2012). Metalliferous and pelagic sediments overlie the UV2 lava and are covered by the V3 lava. Based on our field examination for several sections in the Wadi Hilti area, the stratigraphy of the sediments on the UV2 lava consists of metalliferous sediments, red mudstone and micritic limestone, in ascending order. We obtained *Guttacapsa gutta* and *Thanarla pulchra* from red mudstone. According to O'Dogherty (1994), the co-occurrence of these species is restricted to be middle to late Cenomanian. From red mudstone and micritic limestone, *Rhopalosyringium scissum* and *Hemicryptocapsa polyhedra* were recovered, which can be assigned to a Turonian age (O'Dogherty, 1994).

Very recently, high-precision U-Pb zircon datings were conducted by Rioux et al. (2012, 2013, 2014) on gabbros, tonalites, and trondhjemites. According to them, the rocks fall into two groups in terms of their structural positions, dates, and isotopic composition: the older group, dated at ca. 96.5–95.5 Ma, is attributed to ridge magmatism (V1 lava), whereas the younger group (ca. 95.5–95.0 Ma) is related to post-ridge magmatism (V2 lava). Hara and Kurihara (2015) reported the maximum age of sediments on the V1 lava to middle–late Cenomanian. In the present study, the sediments on the UV2 lava can be correlated with the middle to late Cenomanian. These findings reveal that the activity from the V1 to UV2 lavas was terminated in late Cenomanian. In addition, these radiolarian ages are consistent with the high-precision U-Pb zircon ages of crustal rocks formed by ridge and post-ridge magmatisms. These age constraints imply that the change of tectonic setting progressed rapidly in a short period of middle–late Cenomanian.

Keywords: Oman Ophiolite, radiolarians, Cretaceous, Cenomanian

Radiolarian biostratigraphy of Late Cretaceous pelagic sediments in the Wadi Jizzi area of the Oman Ophiolite

*Kousuke Hara¹, Toshiyuki Kurihara¹

1. Graduate School of Science and Technology, Niigata University

The Cenomanian–Coniacian radiolarian biostratigraphy of pelagic sediments overlying basaltic extrusive lavas of the Oman Ophiolite was studied in the Wadi Jizzi area, about 40 km west of Sohar, northern Oman Mountains. Pelagic and metalliferous sediments commonly occur on lavas and at the boundaries between different volcanic units. The most thickly accumulated sediments, which rest directly on the V1 lava formed by ridge magmatism, were named the Suhaylah Formation (Fleet and Robertson, 1980; Woodcock and Robertson, 1982). The Suhaylah Formation is overlain by the Zabyat Formation (Woodcock and Robertson, 1982; Robertson and Woodcock, 1983). This formation consists of ophiolite debris, redeposited sandstone- to siltstone-sized volcanoclastic rocks, and pelagic mudstone. From the occurrence patterns and stratigraphic ranges of radiolarians, the species clearly make up three distinct assemblages. Based on the occurrences, we defined the following three biostratigraphic zones (interval zones): *Guttacapsa gutta* zone (basal part of the Suhaylah Formation; middle Cenomanian to latest Cenomanian), *Rhopalosyringium scissum* zone (main part of the Suhaylah Formation; latest Cenomanian to Turonian), and *Archaeospongoprimum bipartitum* (Zabyat Formation; Coniacian). In previous studies (e.g., Tippit et al., 1981), the Suhaylah Formation was dated as early Cenomanian to Coniacian–Santonian. We revised the age of this formation to middle–late Cenomanian to Turonian. The radiolarian age of the sediments overlying the V1 lava (ca. 96 Ma) is consistent with the high-precision U–Pb zircon age of crustal rocks formed by ridge magmatism (96.0–95.5 Ma) (Rioux et al., 2014).

Keywords: Oman Ophiolite, radiolarians, biostratigraphy, Cretaceous

Ecology and diversity of phaeodarians (unicellular zooplankton) around Japan

*Yasuhide Nakamura¹, Rei Somiya², Akihiro Tuji³, Noritoshi Suzuki⁴, Rie Hori, S.⁵

1.Hokkaido Univ., 2.Nagasaki Univ., 3.NMNS, 4.Tohoku Univ., 5.Ehime Univ.

Phaeodarians are a group of unicellular zooplankton dwelling in pelagic ocean from the at least Triassic to the present. Phaeodaria have long treated as members of "Radiolaria", however molecular studies revealed that the group in question belongs to Cercozoa. Phaeodarians occasionally become abundant in ocean, and they are thought to have important roles in the marine food web and the material cycles. We will present the results of our investigations concerning this group during the last five years and discuss about the importance of phaeodarians.

Plankton were sampled from several depths at ca. 40 stations around Japan during 2011-2015. The zooplankton in the samples were sorted and identified under a stereomicroscope in order to clarify the composition of each sample. The 18S rDNA sequences of phaeodarians were determined by single-cell PCR method. Some phaeodarians caught in the East China Sea were cultured to observe their behavior.

The microscopic and genetic analyses revealed that two undescribed phaeodarians live in the deep waters in the Sea of Japan, and they were described as *Aulographis japonica* and *Aulosцена pleuroclada*. The former species was abundant through the year, occupying ca. 22% of the total zooplankton biomass on average. The abundance of phaeodarians was also seen in the Kuroshio region, where two species, Aulosphaeridae sp. 1 and Sagosphaeridae sp. 1, occupying 10.2-13.9% of the zooplankton biomass. Another unicellular zooplankton, *Thalassothamnus* sp. 1, was also found in this region. This genus is classified as Entactinaria (Radiolaria) in the current classification system. Our molecular and morphological analyses, however, revealed that the present species is a member of Phaeodaria. Thus, phaeodarians can occasionally become abundant around Japan, but their information is still limited. Further research on unicellular zooplankton near the Japanese Archipelago is necessary for considering the evolution of unicellular organisms in the pelagic ocean.

Keywords: Cercozoa, Phaeodaria, plankton, present, protist, silica

Parameter identification of mathematical models based on three dimensional information of planktic foraminiferal shell structure obtained by Microfocus X-ray CT

*Naoko Kishimoto¹, Katsunori Kimoto²

1.Setsunan University, 2.Japan Agency for Marine-Earth Science and Technology

Morphological variations in microfossils like foraminifers and radiolarians might have the meaning of various functions for their evolution and adaptation in the geological history. In order to quantitatively understand such morphological variations, it is important to parameterize the morphological features in microfossils. We obtained precise morphology of a modern planktic foraminifera *Globigerinoides ruber* (d'Orbigny) using Micro-Xray Computed Tomography (Micro-XCT) technique, and developed numerical approximate models to validate the parameterization by experimental rules used the sequentially connected spheres. Our approximate spherical model clarified that several parameters including radius ratio, distances of geometric centers in each chambers are almost constant, and others such as revolving angle between adjacent chambers are gradually transformed associated with growth of *G. ruber*. It indicates that previous numerical model could not interpret whole morphology of this species. The coupling of mathematical model and Micro-XCT technique has the great potential to realize various forms of microfossils and understand its functional morphology.

Keywords: Microfocus X-ray CT, Mathematical Model, functional morphology, Planktic foraminifera

A search for impact ejecta deposits in the Upper Triassic limestone of the Pizzo Mondello section, western Sicily, Italy.

*Tatuya Kuranari¹, Katsuhito Soda¹, Daisuke Yamashita¹, Tetsuji Onoue¹

1.Department of Earth and Environmental Sciences, Graduate School of Science and Technology, Kumamoto University

The 34-million-year (My) interval of the Late Triassic is marked by the formation of several large impact structures on Earth, including the 90-km-diameter Manicouagan impact crater in Quebec, Canada. This crater occurred in the middle-upper Norian (ca. 215.5 Mya). Late Triassic impact events have been considered a factor in biotic extinction events in the Late Triassic (e.g., end-Triassic extinction event), but this scenario remains controversial because of a lack of stratigraphic records of ejecta deposits. The Manicouagan impact ejecta deposits are known only from Mino terrane, Central Japan and southwestern Britain. To investigate the impact ejecta deposit derived from the Manicouagan impact crater, we examined the stratigraphic variations in major and trace element concentrations from the Upper Triassic Pizzo Mondello section in western Sicily, Italy. The Pizzo Mondello consists of a continuously exposed sequence (a 450m thick) of upper Carnian through late Norian (Upper Triassic) pelagic limestone. The trace element data from this study show a consistent trend of decreasing Cr and Ni values starting in the lower Norian. Concentration of these elements are observed in the upper part of the Pizzo Mondello section and is roughly located at the base of the *Epigondolella bidentata* conodont zone in the late Norian.

Keywords: Late Triassic, ejecta sediment, Pizzo Mondello, limestone, main trace element analysis

Study on analytical technique of carbon isotopes for samples with minute and lean organic carbon content.

*Keiichiro Hasebe¹, Akiko S. Goto², Takashi Hasegawa²

1.Division of Environmental Science and Engineering, Graduate School of Natural Science and Technology, Kanazawa University, 2.School of Natural System, College of Science and Engineering, Kanazawa University

Stable organic carbon isotope ratio ($\delta^{13}\text{C}_{\text{org}}$) of oceanic sediments is useful to discuss paleoceanographic change (e.g. increasing productivity on ocean surface, the global carbon cycle). However, very low total organic carbon content (TOC) makes carbon isotope analysis of organic carbon with regular equipment quite difficult. A shortage of stratigraphic record of $\delta^{13}\text{C}$ from pelagic sediments including chert is partly owe to it.

This study tried carbon isotope measurement of minuscule quantity of carbon and low TOC samples with techniques of elementary analyzer introduction interface (EA/IRMS) and dual inlet. EA/IRMS is online method that uses continuous-flow system and dual inlet is off-line method that is an analysis with IRMS of dual inlet type after cryogenic purification with a glass vacuum line. L-alanine were mainly used for experiments of EA/IRMS method as a standard material. For dual inlet method, standards (ANU-Sucrose, L-alanine, eicosane, triphenylamine) were mostly used. A few chert samples were used for each experiment as a natural sample. Experimental contents of EA/IRMS method were: 1) to know how minute the sample is enough to analyze TOC with elemental analyzer (EA), 2) to know how much weight of powdered sample can be introduced for the measurement of $\delta^{13}\text{C}_{\text{org}}$ and 3) to test actual organic carbon isotope analysis with chert sample. They demonstrated that EA/IRMS method can deal with carbon as much as 10 μg , one-fifth of suitable carbon weight for the conventional analysis in our lab. In spite of it, we judged that EA/IRMS method did not suit for measurement of chert as the volume of chert powder was too much to introduce to the equipment, as an analysis effected on the next analysis (memory effect), and as it damaged the analytical device. For dual inlet method, 1) estimation how minute amount of CO_2 gas is able for the analysis, 2) evaluation the stability of $\delta^{13}\text{C}_{\text{org}}$ for minuscule samples, 3) evaluation of calibration line by multiple standard samples, 4) preliminary organic carbon isotope analysis of actual chert sample, were conducted in this study. The results showed that the values obtained with 50 μgC standards were similar to that with 500 μgC standards (the amount for regular analysis). Moreover, dual inlet method demonstrated that $\delta^{13}\text{C}_{\text{org}}$ value from chert (~700 mg) were not stabilized (showed >1% difference) in replicated sample. On the other hand, the difference of $\delta^{13}\text{C}_{\text{org}}$ between 760 mg of diluted L-alanine (with baked quartz sand) controlled as low TOC (0.007%) and non-diluted L-alanine 0.1mg was about 0.5% showing more stable result than chert samples. Although analysis of standard samples with 50 μgC and low TOC condition with dual inlet method is nearly achieved, problems related to specific character of chert may be remained unresolved for organic carbon isotope analysis of chert.

Keywords: organic carbon isotope ratio, low total organic carbon content (TOC), dual inlet, Elemental Analyzer/ Isotope Ratio Mass Spectrometer (EA/IRMS)

Reconstruction of seamounts by stratigraphy and geochemistry of greenstones in an accretionary complex in the Mitsuishi area, Hokkaido, Japan

*Megumi Ozawa¹

1. Graduate School of Science & Technology, Niigata University

Seamounts and oceanic islands are ones of the common volcanoes on the earth. Their structures and growth processes have been modeled mainly by observation of deep sea drill cores and uplifted submarine volcanic complex (ex. Schmincke, 2010). However, modern volcanoes can provide limited information on their internal structure.

A large amount of greenstones occurs in the Kamuikotan zone of Hokkaido, Japan. Nakano&Komatsu (1979) interpreted that they originated from oceanic island alkali basalts based on the common occurrences of aegirine and kaersutite. While Kimura et al. (1994) assumed an oceanic plateau for their origin based on geochemistry of greenstones, plate tectonic reconstruction, and occurrences of chert xenoliths. Sakakibara et al. (1999) considered its age as Late Jurassic based on radiolarian fossils from chert xenoliths. Although their origins are still controversial, eruptive structures and stratigraphy preserved in the southern Kamuikotan zone will provide further knowledge on internal structure of intra-plate submarine volcanoes.

Therefore, this presentation tries to reconstruct volcanic structures from stratigraphy, eruptive and sedimentary features, and geochemistry of greenstones in the Kamuikotan zone of the Mitsuishi area.

The Futagawa unit along the Futagawa River, a tributary of the Mitsuishi River, is characterized by non-deformed greenstones associated with sedimentary rocks, and shear zones consisting mainly of pelitic mixed rock.

Greenstones in the Futagawa unit can be classified into slightly enriched mid-ocean ridge basalt (T-MORB), ocean island alkali basalt (OIA), and ocean island basalts showing a transitional composition of tholeiitic and alkaline series (OITB). The Futagawa unit is subdivided into sheets A, B, and C, bounded by the shear zones.

Sheet A consists of pillow lava of T-MORB and OITB, dolerite sills of OIA, and sedimentary rocks such as chert and clastic limestone at the bottom of the sequence. Therefore, it is speculated to have formed at a foothill of a volcano in the deep sea during the initial undersea volcano stage of ocean island formation. Sheet B consists of chert at the bottom, overlying pillow lava of relatively homogenous OITB, and large amounts of dolerite sill. It was considered to have formed at a deep sea foothill of the volcano in the shield volcano stage. Sheet C consists of vesicular lavas and pyroclastic rocks of OITB, with a dike crosscutting them. It suggests a central part of the undersea volcano at a relatively shallow level, although it is difficult to assign exact evolutionally stage.

Keywords: seamount, accretionary complex, greenstone

Detrital clinopyroxene and zircon in the Tokoro Belt, Hokkaido: Insight to Cretaceous magmatic arc in the Sea of Okhotsk

*Hayato Ueda¹, Yuta Akutsu¹

1.Department of Geology, Niigata University

Basement of the Sea of Okhotsk consist of ~20 km thick continent-like crust. Jurassic and younger granites and volcanic rocks have been dredged from its sea floor. It is thus regarded as a continental block (Okhotsk Sea Block: OSB). However, rocks suggestive of Precambrian basement and Paleozoic strata common in continental blocks in NE Asia derived from the super-continent have not been found from the sea. Therefore, the origin of OSB is left unknown and could be different from the common continental blocks. It is however difficult to directly explore the OSB basement in detail because it is mostly submerged in the sea. Cretaceous and Paleogene in the Nemuro and Tokoro belts of eastern Hokkaido belong to the paleo-Kuril arc of the OSB margin. They are characterized by volcanogenic sandstones, whose clastic sources are assumed to be a volcanic arc in OSB. They can thus be a key to understand age and composition of OSB. This paper tests chemical composition of clinopyroxene and ages of zircons in sandstones of the Tokoro Belt. Detrital clinopyroxenes are common in sandstones of the Nikoro Group (middle? Cretaceous accretionary complex), Saroma Group (Late Cretaceous forearc basin), and the Yubetsu Group (early Paleogene accretionary complex). All of analyzed clinopyroxene grains are poor in titanium and resemble to those of island arc volcanics. The Nikoro and Saroma sandstones contain scarce and tiny zircon crystals. Among them, grains with enough sizes for LA-ICPMS measurement yielded approx. 90-100 Ma. On the contrary, the Yubetsu sandstone yielded abundant zircon grains, whose U-Pb ages cluster in the Late Cretaceous (65-80 Ma), Jurassic, Permian, and Paleoproterozoic to Neoarchean. A large age gap exists between 430 and 1600 Ma in the sample. Occurrences of detrital zircons suggest that a distinct change in clastic supply occurred in the end of Cretaceous. Detrital zircons in the Yubetsu sandstone share the common age distribution with those in the NE Japan arc, whereas the Yubetsu Group still has characteristics of paleo-Kuril arc with abundant intermediate to mafic volcanic clasts and clinopyroxene composition common with the Nikoro and Saroma groups. Such coexistence of both NE Japan (Eurasia) and Kuril (Okhotsk) natures in the Yubetsu Group is consistent with the dual provenance model by Nanayama (1992), in which the group deposited at the junction between the two arc-trench systems. It is reasonable to consider that the two arcs joined in late Late Cretaceous, so far as no sign of clastic supply from the Eurasia is found in the Saroma and Nikoro groups. These groups formed before the coupling might be more important to examine the origin of OSB using clastic composition.

Keywords: Tokoro Belt, Okhotsk Sea, Paleo-Kuril arc

Sandstone dikes within pelagic cherts of the Permian Yoshii Group in Southwest Japan and its deformation structure

*Tsuyoshi Ito¹, Atsushi Matsuoka²

1.Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, 2.Department of Geology, Faculty of Science, Niigata University

Deformation structure of consolidated cherts would record its diagenetic process. The Akiyoshi terrane, which is a Permian accretionary complex of the Inner Zone of Southwest Japan, contains Permian cherts. However, there are a few studies that have described the deformation structure of the Permian cherts of the Akiyoshi terrane. The Yoshii Group, distributed over west Okayama Prefecture, belongs to the Akiyoshi terrane.

We discovered ductilely- and brittlely-deformed chert blocks within sandstone dikes of the Yoshii Group in the Mihara area. The ductilely-deformed chert blocks and the clay minerals within the sandstone dikes are elongated and have slightly preferred orientation.

Pseudoalbaillella sp. cf. *P. longtanensis* Sheng et Wang was obtained from a chert bed just below the sandstone dike. *Pseudoalbaillella longtanensis* is characteristic species of the *P. longtanensis* Assemblage-Zone corresponding tentatively to the Kungurian (Cisuralian: Lower Permian).

The presence of the ductilely-deformed cherts suggests that the siliceous deposits had yet been unconsolidated when the sandstone dikes intruded. The sandstone dikes might be originated in terrigenous clastic, which is a component of chert-clastic sequences of the Yoshii Group in the Mihara area. The Permian chert had kept unconsolidated more than 10 My in this case.

Keywords: ductile deformation, Permian, radiolaria, pelagic chert, sandstone dike, Akiyoshi terrane

InterRad 15 in Japan (October, 2017) and outreach activities for radiolarians

*Atsushi Matsuoka¹

1.Department of Geology, Faculty of Science, Niigata University

InterRad is the international association of radiolarists. The 15th InterRad meeting will be held in Japan during 23-27 October in 2017. In relation to this international meeting, we are planning to do something in enhancing a degree of recognition of radiolarians. Our projects include not only research and education but also outreach activities. It is essential for any academic associations to introduce relevant topics into society. The organizing committee of the InterRad 15 will carry out outreach activities continuously and systematically by using available tools.

Keywords: Radiolaria, InterRad, outreach