

マレーシアサラワク州の3つの熱帯泥炭地における土壤CO<sub>2</sub>およびCH<sub>4</sub>フラックス観測の初期結果  
Initial results of observations of soil CO<sub>2</sub> and CH<sub>4</sub> fluxes in three ecosystem types of tropical peatland in Sarawak, Malaysia

\*平田 竜一<sup>1</sup>、沖元 洋介<sup>2</sup>、平野 高司<sup>2</sup>、Kiew Frankie<sup>2,3</sup>、Melling Lulie<sup>3</sup>

\*Ryuichi Hirata<sup>1</sup>, Yosuke Okimoto<sup>2</sup>, Takashi Hirano<sup>2</sup>, Frankie Kiew<sup>2,3</sup>, Lulie Melling<sup>3</sup>

1.国立環境研究所、2.北海道大学 農学研究院、3.Tropical Peat Research Laboratory Unit

1.National Institute for Environmental Studies, 2.Research Faculty of Agriculture, Hokkaido University, 3.Tropical Peat Research Laboratory Unit

Tropical peatlands in Southeast Asia store large carbon by accumulating peat and they are vulnerable to climate change and human disturbances.

Soil carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>) fluxes observations were started on September 2015 at three ecosystem types of tropical peatland in Sarawak, Malaysia. The sites were one tropical swamp forest with high ground water level and one with low ground water level, and an oil palm plantation on peat. In each site, we installed an automated multi-chamber system. We will present the initial results of the observations.

キーワード：自動チャンバーシステム、泥炭林、オイルパーム

Keywords: Automated multi-chamber system, peat swamp forest, oil palm