

大規模攪乱後の植生遷移初期におけるCO₂フラックスの変化Change of CO₂ flux during an early secondary succession after severe forest disturbance*平野 高司¹*Takashi Hirano¹

1.北海道大学大学院農学研究院

1. Research Faculty of Agriculture, Hokkaido University

Forest ecosystems are expected as a major sink of atmospheric carbon dioxide (CO₂), whereas their ability to absorb CO₂ is severely perturbed by disturbances, such as deforestation, fires, windthrow etc. Many studies which adopted the chrono-sequence approach reported that such severe disturbances often changed forest to be a CO₂ source. However, there are few studies that directly measure CO₂ flux for a long term (more than 10 years) above a disturbed forest site during an early stage of secondary succession after severe disturbance. A flux site of a larch plantation in Tomakomai, Hokkaido, Japan was struck by a typhoon in September 2004. Because of wind storm, about 90% of trees fell down. The fallen trees were removed by heavy machinery from the site, through which the soil surface and understory species were also disturbed. After the operations of timber transport, secondary succession progressed naturally in the ex-forest site. We recommenced flux measurement in August 2005. CO₂ flux has been measured by the eddy covariance technique with an open-path CO₂ / H₂O analyzer (LI7500, Licor) during a snow-free period from mid-April to mid-November. Cumulative net ecosystem CO₂ exchange (NEE) during the snow-free period was positive every year until 2015, whereas it showed a negative relationship, which indicates that the CO₂ source strength of the ecosystem decreased. This negative relationship was caused by the increase of gross primary production (GPP) or ecosystem photosynthesis, which corresponded to vegetation recovery through secondary succession.

キーワード：風倒害、渦相関法、植生回復

Keywords: Windthrow, Eddy covariance, Vegetation recovery