

## グリーンランド北西部における溢流氷河の表面高度変化

## Surface elevation change of outlet glaciers in northwestern Greenland

\*津滝 俊<sup>1,2</sup>、片山 直紀<sup>2,3</sup>、榊原 大貴<sup>2,3</sup>、杉山 慎<sup>2</sup>、澤柿 教伸<sup>4</sup>

\*Shun Tsutaki<sup>1,2</sup>, Naoki Katayama<sup>2,3</sup>, Daiki Sakakibara<sup>2,3</sup>, Shin Sugiyama<sup>2</sup>, Takanobu Sawagaki<sup>4</sup>

1.国立極地研究所国際北極環境研究センター、2.北海道大学低温科学研究所、3.北海道大学大学院環境科学  
院、4.法政大学社会学部

1.Arctic Environment Research Center, National Institute of Polar Research, 2.Institute of Low  
Temperature Science, Hokkaido University, 3.Graduate School of Environmental Science, Hokkaido  
University, 4.Faculty of Social Science, Hosei University

Ice discharge from calving glaciers has been increasing in the Greenland ice sheet (GrIS) since 2000s. This increase plays important roles in the volume change of GrIS and its contribution to sea level rise. To investigate the mass loss of GrIS calving glaciers, ice surface elevation change has been studied by differencing digital elevation models (DEMs) derived by satellite remote-sensing. Panchromatic Remote-sensing Instrument for Stereo Mapping (PRISM) images of Advanced Land Observing Satellite (ALOS) have a spatial resolution of 2.5 m, which is fine enough to measure several meters of elevation change on glaciers. The large spatial coverage of the images (1225 km<sup>2</sup>) is suitable for studying glaciers distributed over a large area.

In this study, we measured surface elevation change of 14 outlet glaciers near Inglefield Bredning in northwestern Greenland (77°47′–78°10′N, 65°00′–72°47′W). These glaciers flow into the ocean except for two land-terminating glaciers. We processed stereo pair ALOS PRISM images acquired in summer 2007 and 2010 with a digital map plotting instrument (Leica Photogrammetry Suite) to generate DEMs with a 25 m grid mesh. Elevation data from ALOS DEMs were calibrated on ice-free terrain, and compared to calculate ice surface elevation change between 2007 and 2010.

The surface elevation of all the studied glaciers decreased and the magnitude of the elevation change increases downglacier. The mean elevation change rate ranged from -0.4 to -4.9 m a<sup>-1</sup>. Marine-terminating Tracy and Bowdoin Glaciers thinned at rates of -4.9 and -4.1 m a<sup>-1</sup>, which were larger than those at other glaciers. The rate at Tugto Glacier, a land-terminating glacier located near Bowdoin Glacier, was -2.8 m a<sup>-1</sup>. This result confirms that recent thinning of GrIS outlet glaciers is more significant at marine-terminating glaciers as compared to land-terminating glaciers. Rapid thinning of marine-terminating outlet glaciers observed in this study suggests the importance of ice dynamics and/or ice-ocean interaction in the mass loss of GrIS.

キーワード：北極圏雪氷学、氷河表面高度変化、氷河ダイナミクス

Keywords: Arctic glaciology, Glacier thinning, Glacier dynamics