

バイカル湖天然ガスハイドレートの水和数の圧力依存性

Effect of water depth on hydration number of natural gas hydrate in Lake Baikal

*太田 有香¹、八久保 晶弘¹、Khlystov Oleg²、Kalmychkov Gennadiy³、De Batist Marc⁴、坂上 寛敏¹、南 尚嗣¹、山下 聡¹、高橋 信夫¹、庄子 仁¹

*Yuka Oota¹, Akihiro Hachikubo¹, Oleg Khlystov², Gennadiy Kalmychkov³, Marc De Batist⁴, Hirotohi Sakagami¹, Hirotsugu Minami¹, Satoshi Yamashita¹, Nobuo Takahashi¹, Hitoshi Shoji¹

1.北見工業大学、2.ロシア科学アカデミー陸水学研究所、3.ロシア科学アカデミー地球化学研究所、4.ゲント大学

1.Kitami Institute of Technology, 2.Limnological Institute, SB RAS, 3.Vinogradov Institute of Geochemistry, SB RAS, 4.Ghent University

Natural gas hydrates exist under the deep sea/lake or permafrost are considered to be a potential natural gas resource. Hydration number "n" of methane hydrate ($\text{CH}_4 \cdot n\text{H}_2\text{O}$) decides the amount of methane in an unit volume/weight of crystal. In the case of ideal full-occupation of hydrate cages, the value of hydration number is 5.75 (Sloan and Koh, 2008). However, researchers have reported that the hydration number is around 6, because small amount of empty cages decrease the free energy and stabilize the crystal. Natural gas hydrates have been retrieved from lake-bottom sediment at Lake Baikal, where the water depths ranged from 450m to 1400m, and their main gas component is methane. Hydration number may change under various pressure condition, but it has not been examined yet. In this study, we report the pressure effect on hydration number of synthetic methane hydrate and natural gas hydrate of Lake Baikal.

Methane hydrate was synthesized under the pressure range between 3 MPa to 20 MPa. Natural hydrate samples were retrieved at the southern Baikal basin (Malenky, Bolshoy, Peschanka P-2, and Goloustnoye G-1) and central Baikal basin (Kukuy K-1, K-2, K-8, K-9, K-10, and Novosibirsk). Raman spectroscopic measurements were made to assess the hydration numbers of samples. Raman spectra were obtained at 123 K in the range 2,800-3,000 cm^{-1} for the C-H stretching peaks of methane, and fitted using a Voigt function to obtain the integrated intensities of the two peaks corresponding to methane engaged in large and small cages of the cubic structure I. The cage occupancies and the hydration numbers were estimated from these peak intensities using a statistical thermodynamic model (Sum et al., 1997). Hydration number of synthetic methane hydrate decreased with pressure, from 6.05 (2.7 MPa) to 5.97 (20.9 MPa), and those of natural gas hydrate also decreased slightly with water depth.

We are grateful to the support of the crew onboard R/V Vereshchagin during the cruises in Lake Baikal (MHP project). This work was supported by funding agencies in Japan (Grant-in-Aid for Scientific Research 24404026 and 26303021 from the Japan Society for the Promotion of Science). Sloan and Koh (2008) Clathrate Hydrates of Natural Gases, 3rd ed., CRC Press: Boca Raton, FL, USA Sum et al. (1997) Measurement of clathrate hydrates via Raman spectroscopy. J Phys Chem B 101: 7371-7377.

キーワード：ハイドレート、水和数、バイカル湖、ラマン分光分析

Keywords: hydrate, hydration number, Lake Baikal, Raman spectroscopic analysis