

2015年3月14日に観測されたディスクリートアークのオーラトモグラフィ解析初期結果

Preliminary results of auroral tomography analysis of discrete arcs observed on March 14, 2015

*田中 良昌¹、小川 泰信¹、門倉 昭¹、Gustavsson Bjorn²、Partamies Noora³、Kauristie Kirsti⁴、Whiter Daniel⁵、Enell Carl-fredrik⁶、Braendstroem Urban⁷、Sergienko Tima⁷、Kozlovsky Alexander⁸、宮岡 宏¹、Kosch Mike⁹

*Yoshimasa Tanaka¹、Yasunobu Ogawa¹、Akira Kadokura¹、Bjorn Gustavsson²、Noora Partamies³、Kirsti Kauristie⁴、Daniel Whiter⁵、Carl-fredrik Enell⁶、Urban Braendstroem⁷、Tima Sergienko⁷、Alexander Kozlovsky⁸、Hiroshi Miyaoka¹、Mike Kosch⁹

1.国立極地研究所、2.UiT The Arctic University of Norway, Norway、3.The University Centre in Svalbard, Norway、4.Finnish Meteorological Institute, Finland、5.University of Southampton, UK、6.EISCAT Scientific Association、7.Swedish Institute of Space Physics, Sweden、8.Sodankyla Geophysical Observatory, Finland、9.South African National Space Agency, South Africa

1.National Institute of Polar Research, 2.UiT The Arctic University of Norway, Norway, 3.The University Centre in Svalbard, Norway, 4.Finnish Meteorological Institute, Finland, 5.University of Southampton, UK, 6.EISCAT Scientific Association, 7.Swedish Institute of Space Physics, Sweden, 8.Sodankyla Geophysical Observatory, Finland, 9.South African National Space Agency, South Africa

We conducted a campaign of auroral tomography observation using multi-point imager network and the EISCAT UHF radar in Northern Europe in March, 2015. During this campaign, an auroral breakup was observed at 23:15 UT on March 14 by three all-sky EMCCD imagers and three wide-view CCD imagers. Wavy structure of discrete arcs was often observed around the magnetic zenith at Tromso from 22 to 23 UT and pulsating aurora was observed after the breakup. The monochromatic (427.8 nm) images were taken at a sampling interval of 2 seconds by the three EMCCD imagers and at an interval of 10 seconds by all the six imagers. The EISCAT UHF radar at Tromso measured the ionospheric parameters along the field line at the magnetic zenith from 20 to 24 UT.

We apply the auroral tomography method to these data set to reconstruct reliable three-dimensional distribution of the 427.8 nm emission, that will allow us to investigate quantitatively the following subjects; (1) relation between the 427.8 nm emission and electron density enhancement along the field line, (2) spatial distribution of energy of precipitating electrons in the wavy structure, in particular, relation between the energy of precipitating electrons and thickness of discrete arc, (3) relation between motion of the wavy structure and spatial distribution of the ionospheric conductivity, and (4) spatial and temporal variations of energy distribution of precipitating electrons at the auroral breakup. We present preliminary results from the auroral tomography analysis of the discrete arcs.

キーワード：オーロラ、トモグラフィ解析、3次元分布、電離圏、EISCATレーダー

Keywords: aurora, tomography analysis, 3D distribution, ionosphere, EISCAT radar