Effect of pressure on water solubility in aluminous bridgmanite Effect of pressure on water solubility in aluminous bridgmanite

\*Chen Jiuhua<sup>1,2</sup>, Pamato Martha<sup>3</sup>, Inoue Toru<sup>4</sup>, Kakizawa Sho<sup>4</sup>, Yang Bin<sup>1</sup>, Lin Yangting<sup>5</sup>, Katsura Tomoo<sup>3</sup>, Kawazoe Takaaki<sup>3</sup>, Liu Bingbing<sup>6</sup> \*Jiuhua Chen<sup>1,2</sup>, Martha G Pamato<sup>3</sup>, Toru Inoue<sup>4</sup>, Sho Kakizawa<sup>4</sup>, Bin Yang<sup>1</sup>, Yangting Lin<sup>5</sup>, Tomoo Katsura<sup>3</sup>, Takaaki Kawazoe<sup>3</sup>, Bingbing Liu<sup>6</sup>

 Florida International Univ. 2.Center for High Pressure Science and Technology,
Changchun, 3.Bayerisches Geoinstitut, Universitaet, 4.Ehime University, 5.Institute of Geology and Geophysics, Chinese Academy of Sciences, 6.Jilin University
Florida International Univ., 2.Center for High Pressure Science and Technology, Changchun,
Bayerisches Geoinstitut, Universitaet, 4.Ehime University, 5.Institute of Geology and Geophysics,
Chinese Academy of Sciences, 6.Jilin University

We have conducted a sequence of high pressure experiments to study water solubility in aluminous bridgmanite as a function of pressure at  $1900^{\circ}$ C. The experimental high pressures were generated using multi-anvil presses at Bayerisches Geoinstitut (BGI) for pressures up to 28 GPa and at Geodynamics Research Center (GRC) for pressures above 30GPa. The starting material for these experiments was a mixture of oxides (Mg(OH)<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub> and SiO<sub>2</sub>) with equivalent of about 5 mol % of Al  $_2O_3$  and 15 wt % of H<sub>2</sub>O. The structure and composition of the bridgmanite sample after high P/T syntheses were examined using x-ray diffraction (XRD) and electron probe microanalysis (EPMA). Water concentration in the sample was measured using secondary ion mass spectroscopy (SIMS) and Fourier transform infrared spectroscopy (FTIR). The measurements yield that the aluminous bridgmanite with about 2 wt% of Al<sub>2</sub>O<sub>3</sub> may take as much as 0.13 wt % of H<sub>2</sub>O at the P/T condition of the top of the Earth's lower mantle and this solubility increases significantly with pressure. At the bottom of Earth's lower mantle, bridgmanite may take nearly a couple of weight percent of water reservoir of the lower mantle can be as large as a few to ten oceans of water.