

ローソン石青色片岩の相平衡関係と吸水・脱水機構：黒瀬川構造帯箱石サブユニットの例

Phase relation of lawsonite-blueschists and their role as a water budget: A case study from the Hakoishi sub-unit of the Kurosegawa belt, SW Japan

*平島 崇男¹、佐藤 永太郎¹、吉田 健太¹、上村 謙一郎¹、藤本 善航¹

*Takao Hirajima¹, Eitaro Sato¹, Kenta Yoshida¹, Kenichiro Kamimura¹, Kazuyoshi Fujimoto¹

1.京都大学大学院理学研究科地球惑星科学専攻

1.Department of Geology and Mineralogy, Graduate School of Science, Kyoto University

This study investigated progressive changes in the mineral assemblage of a Palaeozoic lawsonite-blueschist (BS) unit of the Kurosegawa belt, Kyushu, Japan. The BS unit is mainly composed of intercalated metachert and metabasaltic rocks surrounded by serpentinite. Based on the spatial distribution of mineral assemblages in metabasaltic rocks, three mineral zones can be identified in a ca. 10 km-long east-west oriented unit: (1) a pumpellyite (Pmp) + Na-amphibole (Namp) assemblage dominated area (Zone 1); (2) an intermixture area of lawsonite (Lws) + Na-pyroxene (Napx) + Pmp and Lws + Namp + Pmp assemblages (transition zone); and (3) an area dominated by Lws + Namp + Napx assemblages (Zone 2), with excess chlorite (Chl), albite (Ab), quartz (Qz), titanite, and Fe oxides. Na-pyroxene varies its composition from jadeite(Jd)₂₀-diopside(Di)₂₅ of Zone 1 to Jd₃₅Di₁₅ in Zone 2 with a similar composition range of aegirine(Ac) content (from 40 to 55). Glaucophane component {XGln = Al/(Al+Fe³⁺)} of Na-amphibole associated with Lws and Pmp increases from XGln = 0.2 in Zone 1 to XGln = 0.8 in Zone 2. Jd contents and a lack of epidote and zeolite group minerals suggest that the pressure (P) conditions of the BS increase from Zone 1 (0.50 GPa) to Zone 2 (0.75 GPa) between 200-300 °C, suggesting that a low geothermal gradient (ca. 5-10 °C/km). Schreinemaker's analyses in a NaCaMgFe₃AlSiH (NCF3ASH) system using mean mineral compositions of representative samples with excess of Chl, Ab, Qz and H₂O suggest that 1) observed variations in mineral assemblages can be formed by a hydration reaction of Pmp + Napx + Chl + H₂O = Lws + Namp, which was driven by a P increase from Zone 1 to Zone 2 and they represent the transition from the Pmp-BS to Lws-BS sub-facies; 2) the total H₂O content stored in hydrous minerals increases from 3.6 wt % in the Pmp-Namp assemblage of Zone 1 to 5.0-6.4 wt % in the Lws + Namp + Napx assemblages in Zone 2; and 3) along with the Lws-Namp and Pmp-Namp assemblages, the Napx + Chl assemblage can retain a significant volume of H₂O and its stability field can expand to higher P conditions by decreasing the Di and increasing the Jd and Ac contents of Napx.

キーワード：ローソン石青色片岩、相平衡、沈み込み帯流体、黒瀬川構造帯

Keywords: lawsonite blueschist, phase relation, H₂O budget in subduction zone, Kurosegawa belt.