

## Petrologic Study of the Stage IV Eruptives of Myoko Volcano

\*Rin Nodera<sup>1</sup>, Yasuo Ishizaki<sup>1</sup>, Nobuko Kametani<sup>1</sup>

1. Graduate School of Science and Engineering for Education, University of Toyama

Myoko Volcano (2445 m a.s.l.), a near-conical composite stratovolcano with a summit horseshoe-shaped depression ~1.5 km wide, is located at the western Niigata Prefecture. To date, the eruptive history and characteristics of this volcano have well studied. The volcanic activity started at 0.3 Ma and is divided into four eruption stages on the basis of K-Ar and <sup>14</sup>C ages (Hayatsu, 2008; hereafter H08). In order to decipher the magmatic evolution of the composite volcano, petrographic features and whole-rock compositions of the Stage IV eruptives are investigated.

The stage IV, the youngest stage of Myoko Volcano, consists of three sub-stages, i.e., pre-collapse (43 ka-), collapse (21 ka-6 ka), and post-collapse stages (6 ka-present) (stage names are modified from those in H08). The volcanic activity during the pre-collapse sub-stage formed the dacitic to andesitic Shibutamigawa Pyroclastic Flow Deposit (SPFD; 55.2-62.9 wt.% SiO<sub>2</sub>; 42190 ±380 yr BP) and the directly overlying, basaltic Nishikawadani Scoria Flow Deposit (NSFD; 50.5-52.8 wt.% SiO<sub>2</sub>). (<sup>14</sup>C age is from 08). The SPFD shows a hydrous phenocryst assemblage (Pl + Amp + Opx + Cpx ± Ol ±Opq) whereas the NSFD shows an anhydrous phenocryst assemblage (Pl ±Opx + Cpx + Ol ±Opq). The volcanic activity during the central cone stage formed two andesitic to dacitic pyroclastic flow deposits, i.e., the Akakura Pyroclastic Flow Deposit (APFD; 5510 ±70 yr. BP, 60.3-64.4 wt.% SiO<sub>2</sub>) and the Ohtagirigawa Pyroclastic Flow Deposit (OPFD; 4060 ±60 yr. BP, 56.9-64.1 wt.% SiO<sub>2</sub>) (<sup>14</sup>C ages are from H08). The APFD and OPFD show hydrous phenocryst assemblage (Pl + Amp + Opx ±Cpx ±Ol ±Qtz ±Opq) and contain quenched mafic enclaves (55.2-57.1 wt.% SiO<sub>2</sub>; phenocryst assemblage Pl ±Amp + Opx + Cpx ±Ol ±Opq). Eruptives of pre- and post-collapse sub-stages show distinct linear mixing trends on many Harker diagrams. Although major and trace elements are broadly similar in all dacites erupted between pre- and post-collapse stages, there are slight differences in K<sub>2</sub>O contents. In addition, the mafic magmas erupted between pre- and post-collapse stages are differ in TiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub>, K<sub>2</sub>O, P<sub>2</sub>O<sub>5</sub>, V, and Sr contents. These petrological features indicate that the eruptions of the pre- and post-collapse sub-stages were fed by different and transitory silicic magma chambers, repeatedly recharged by mafic magmas of diverse compositions.

Keywords: Myoko Volcano, magma mixing, magma system