Df-P001 Room: Poster Time: June 8 17:30-19:30

Crustal structure of the passive continental margin, west off Svalbard Islands, deduced from ocean bottom seismographic studies

Asako Kuwano [1], Yoshio Murai [2], Yuichi Nishimura [3], Rolf Mjelde [4], Hideki Shimamura [2]

[1] ISV, Hokkaido Univ., [2] Institute of Seismology and Volcanology, Hokkaido Univ., [3] Inst. Seismology and Volcanology, Hokkaido Univ., [4] Institute of Solid Earth Physics, Univ. of Bergen

In 1998, we made controlled source surveys by using the ocean bottom seismographs, airguns and single-channel streamer in the Atlantic Ocean west off Svalbard Islands. We analyzed 3 profiles across the Knipovich Ridge and the passive continental margin in this study. We constructed P-wave velocity structures by ray-tracing method. We can obtain the detailed crustal structure of the ocean-continent transition zone.

In 1998, we made controlled source surveys by using the ocean bottom seismographs, airguns and single-channel streamer in the Atlantic Ocean west off Svalbard Islands. We analyzed 3 profiles across the Knipovich Ridge and the passive continental margin in this study. We constructed P-wave velocity structures by ray-tracing method. We can obtain the detailed crustal structure of the ocean-continent transition zone.