

## Quasi Periodic structures of visible auroras and Syowa East HF radar echoes

# Yozo Murata [1], Natsuo Sato [2], Hisao Yamagishi [3], Akira Sessai Yukimatu [4], Masayuki Kikuchi [2], Kazuo Makita [5], Huigen Yang [2], Ruiyuan Liu [6]

[1] Department of Polar Sci., the Graduate Univ. for Advanced Studies, [2] NIPR, [3] Upper Atmos. Phys., Natl. Inst. Polar Res., [4] UAP, NIPR, [5] Engineering, Takushoku Univ., [6] PRIC

The geomagnetic and geographic latitudes at Zhongshan Station are  $\sim 74.5$  and  $\sim 69.4$  degrees, respectively. The field of views of Syowa East HF Radar covers over Zhongshan Station. Periodic structure of HF radar echoes with period of  $\sim 5$ -10 minutes were found in the period of 12:10 (UT) to 13:50 (UT) in 3 August 1997. Fortunately interesting visible auroras were observed simultaneously using all-sky TV cameras. One of the most interesting characteristics of visible auroras is quasi-periodic variations in the intensity and / or spatial variation of auroras. We examined in detail the relation between the visible auroras and HF radar echoes.

The geomagnetic and geographic latitudes at Zhongshan Station are  $\sim 74.5$  and  $\sim 69.4$  degrees, respectively. The field of views of Syowa East HF Radar covers over Zhongshan Station. Periodic structure of HF radar echoes with period of  $\sim 5$ -10 minutes were found in the period of 12:10 (UT) to 13:50 (UT) in 3 August 1997. Fortunately interesting visible auroras were observed simultaneously using all-sky TV cameras. One of the most interesting characteristics of visible auroras is quasi-periodic variations in the intensity and / or spatial variation of auroras. We examined in detail the relation between the visible auroras and HF radar echoes.