

エジプト東部砂漠のパン・アフリカン（原生代）クロミタイトおよび超マフィック岩類

Pan African (Proterozoic) chromite deposits and associated ultramafic rocks in the Eastern Desert of Egypt

アハメド モハメド [1], 荒井 章司 [1]

Ahmed Hassan Ahmed Mohamed [1], Shoji Arai [2]

[1] 金沢大・理・地球

[1] Dept. Earth Sci., Kanazawa Univ., [2] Dept. Earth Sci., Kanazawa Univ.

<http://kgeopp6.s.kanazawa-u.ac.jp/Arai/Ahmed/>

エジプト東部砂漠のパン・アフリカン・オフィオライトのクロミタイトおよび随伴する岩石の性質を、残留するクロムスピネルの組成により推定した。オフィオライトは著しく分断されメランジュとなり、岩石は変成され、かんらん石は残留しない。クロミタイトは場所により、変成の程度、化学組成、スピネル中の珪酸塩包有物の有無・種類などの変化が大きい。クロムスピネルの $Cr\# (= Cr / (Cr + Al) \text{ 原子比})$ はクロミタイトは比較的高く、0.8に及ぶが、かんらん岩（ダナイト、ハルツバーガイト）では0.5前後である。この予察的データからは、この原生代のオフィオライトの岩石学的性質は基本的に顕生代のものと大差ない。

There are numerous mafic-ultramafic masses of Pan African age in NE Africa and Saudi Arabia that lie within several, broad, discontinuous NW-SE belts. In all cases examined, these masses have mechanically external contacts with the Pan-African rocks already described. These ultramafic bodies account for 5.3 % of all Precambrian outcrops in the Eastern Desert of Egypt. Chromite deposits of Egypt are commonly hosted by serpentized ultramafic rocks widely distributed in the central and southern Eastern Desert of Egypt. Many occurrences of chromite deposits are known : a) Central Eastern Desert: Gebel El Lawi, Wadi El Zarka, Wadi Um Huitate, Wadi Bezah, El Barramyia, Gebel El Rabshi, etc., b) Southern Eastern desert; Abu Dahr and El Gallala. The ultramafic rocks outcropping within the studied areas have been heavily altered into serpentinite, talc carbonate and ultramafic schists. Field and petrographic investigations indicate rarely preserve primary textures and silicate minerals of peridotites, except only one locality (Abu Dahr area, south Eastern desert) where we could observe primary textures of the host ultramafic rocks.

The chromite deposits of Egypt in most cases occur mainly as lenticular bodies of variable sizes generally less than 30 m long. Different ore types have been described including massive ore, disseminated ore and grape to pocket ore. The characteristics of these ore deposits varied from one locality to another in the texture, degree of alteration and chemical composition of spinel. Even in the same locality, the massive ore has different chemical character compared with the disseminated ones. The chemistry of chromian spinel is widely varied from locality to locality partly depending on the degree or grade of alteration or metamorphism. Our preliminary data indicate that the $Cr / (Cr + Al)$ atomic ratio of spinel is up to 0.8 in chromitites and around 0.5 in peridotites for the least metamorphosed samples. This characteristic is almost similar to that of the Phanerozoic ophiolites.