Ea-P006 Room: Lounge Time: June 25 17:30-19:00

Relative contribution of ionospheric currents and magnetospheric currents to mid-latitude geomagnetic field disturbance

Toshihiko Iyemori[1], Shinya Nakano[2], Masahiko Takeda[3], Satoru Yamashita[4], Akinori Saito[2], Hiroaki Yugo[2]

[1] WDC-C2 for Geomag., Kyoto Univ., [2] Dept. of Geophysics, Kyoto Univ., [3] Data Analysis Center for Geomag.and Space Mag., Kyoto Univ., [4] Dept. of Geophysics, Kyoto Univ.

The field-aligned currents which flow into or out from the auroral region generate considerable amount of magnetic variation in m iddle latitudes. we made a comparison between the observed ground magnetic variation and the estimated ionospheric current effects from the electric field data obtained by the MU radar and ionospheric model conductance. The results suggest that the ionospheric current effect is relatively small and most of the disturbance fields in mid-latitudes are magnetospheric origin including the effects of the field-aligned currents.

The field-aligned currents which flow into or out from the auroral region generate considerable amount of magnetic variation in middle latitudes. we made a comparison between the observed ground magnetic variation and the estimated ionospheric current effects from the electric field data obtained by the MU radar and ionospheric model conductance. The results suggest that the ionospheric current effect is relatively small and most of the disturbance fields in mid-latitudes are magnetospheric origin including the effects of the field-aligned currents.