

Redistribution of reactive nitrogen in the Arctic lower stratosphere in the winter of 1999-2000

Makoto Koike[1], Yutaka Kondo[2], Nobuyuki Takegawa[3]

[1] EPS, Univ. of Tokyo, [2] STEL, Nagoya Univ, [3] STEL

We measured total reactive nitrogen (NO_y) in the Arctic lower stratosphere from the NASA DC-8 aircraft during the SOLVE-THESEO campaign in the winter of 1999-2000. The NO_y-N₂O correlation was similar between the December and January deployments. During the March deployment NO_y values were slightly, however, systematically higher as compared to those observed during the previous deployments, although a tight correlation between NO_y and N₂O was still seen. Because of the record low temperature, intensive denitrification was observed at 16-21.5 km. Redistribution due to gravitational sedimentation of PSC particles and evaporation at lower altitudes was considered to be responsible to the observed increase in the NO_y level.

We measured total reactive nitrogen (NO_y) in the Arctic lower stratosphere from the NASA DC-8 aircraft during the SOLVE-THESEO campaign in the winter of 1999-2000. The NO_y-N₂O correlation at 5-12.5 km was similar between the December and January deployments. During the March deployment NO_y values were slightly, however, systematically higher as compared to those observed during the previous deployments, although a tight correlation between NO_y and N₂O was still seen. Because of the record low temperature, intensive denitrification was observed at 16-21.5 km from the NASA ER-2 aircraft. Redistribution due to gravitational sedimentation of PSC particles and evaporation at lower altitudes was considered to be responsible to the observed increase in the NO_y level. The 3D model calculation could generally reproduce the observed features only when the denitrification process was included in the model.