

Liquefaction caused by the 2000 Western Tottori Earthquake, western Japan

aiming lin[1], Ken-ichi Kano[2], Tadashi Maruyama[3]

[1] Institute of Geosciences, Shizuoka Univ, [2] Faculty of Sci., Shizuoka Univ., [3] Institute of Geosciences, Shizuoka Univ.

http://www.sci.shizuoka.ac.jp/~geo/Staff/Lin_j.html

The sand boils induced by the 2000 Western Tottori Earthquake (Mj7.3) widely occurred in reclaimed lands on the Yumigahama sand-bar, northwestern Tottori Prefecture, western Japan. It was observed at several locations that the sand boiling formed by liquefaction was continued for at least two days after the earthquake. This type of liquefaction is called (low liquefaction) in this study. The gush height of boiled sands up to 90 cm was directly observed at some locations. The ground of the sand boiled area has subsided 50 cm in maximum. These phenomena will provide important information both for a quantitative evaluation of liquefaction potential of sand deposits and for a seismic hazard evaluation.

The sand boils induced by the 2000 Western Tottori Earthquake (Mj7.3) widely occurred in reclaimed lands on the Yumigahama sand-bar, northwestern Tottori Prefecture, western Japan. It was observed at several locations that the sand boiling formed by liquefaction was continued for at least two days after the earthquake. This type of liquefaction is called (low liquefaction) in this study. The gush height of boiled sands up to 90 cm was directly observed at some locations. The ground of the sand boiled area has subsided 50 cm in maximum. These phenomena will provide important information both for a quantitative evaluation of liquefaction potential of sand deposits and for a seismic hazard evaluation.