

2001年中国昆仑山中部 Ms8.1 地震の地震断層の性状

Surface ruptures produced by the 2001 Ms 8.1 Central Kunlun (China) earthquake

林 愛明[1], 付 碧宏[2]

aiming lin[1], Bihong Fu[2]

[1] 静岡大・理・地球, [2] 静大・理・地球

[1] Institute of Geosciences, Shizuoka Univ, [2] Graduate School of Science and Technology, Shizuoka Univ.

<http://www.ipcs.shizuoka.ac.jp>

The Ms 8.1 Central Kunlun earthquake of 14 November 2001 was triggered by the active Kunlun fault in the Central Kunlun mountain area, northern Tibet. A co-seismic surface rupture zone of a nearly 400 km long, called Kunlun rupture zone, occurred along the western segment of the Kunlun fault. Field investigations show that the surface ruptures are distributed in a zone with width ranging from few to several hundreds of meters, generally from 5 to 50 m. The rupture zone is composed of distinct shear faults, extensional cracks, and mole tracks. The left-lateral offsets are measured by using the surface deformation marker such as present-day glacier, moraine, stream channel, gully, and road, which vary from few tens of cm to 16.3 m, but generally from 4m to 8 m. The maximum displacement up to 16.3 m was observed across a rupture zone of 550 m wide. Both the rupture length and maximum displacement are the largest among the co-seismic surface rupture zones ever reported worldwide. The co-seismic deformation characteristics of surface markers reveal that the earthquake had a purely strike-slip focal mechanism. Both the geological and geomorphological evidence indicates that the geometry of the ruptures is controlled by the pre-existing Kunlun fault. The large amounts of strike-slip and rupture length produced by the earthquake support the hypothesis that the Kunlun fault plays an important role of strike-slip partitioning in the rapid eastward extrusion of Tibet to accommodate the continuing penetration of Indian plate into Euro-Asian plate. The Central Kunlun earthquake provides an exceptional opportunity to study the geometry of co-seismic rupture structures along a large strike-slip fault for further understanding the relationship between the fault geometry and rupturing process.

2001年11月14日午後5時26分に、チベット高原北部にある昆崙山中部で Ms 8.1 の地震が発生し、中国北西部の広い範囲に強い揺れをもたらした。地震直後の現地調査により、1) この地震に伴い、既存の東西走向を持つ昆崙断層沿いに長さ 400 km 以上におよぶ地表地震断層が現れたこと、2) 地震断層はほぼ純粋な左横ずれであること、3) 左横ずれ量は最大で 16.3 m に達すること、4) 地震断層の割れ目帯の幅は数メートルから 10 km まで変化することなどが、観察された。この地表地震断層帯長さと変位量はこれまでに世界中に報告されたものの中でもっとも大きい。この地震断層の長さと横ずれ変位量の大きさは、昆崙断層がインド・プレートとユーラシアプレートの衝突によるチベット高原の東への extrusion 過程で横ずれ partitioning の役割を果たしていることを示した。