

## Seismo-Atmospheric Disturbances Observed by Anomalous Propagation of VHF Electromagnetic Waves

Hironobu Fujiwara[1], # Masashi Kamogawa[2], Jann-Yenq Liu[3], Yu-Jung Chuo[4], Hideho Ofuruton[5], Yoshi-Hiko Ohtsuki[1]

[1] Dep. of Phys., Waseda Univ., [2] Dep. of Phys., Tokyo Gakugei Univ., [3] Inst. of Space Sci., NCU in Taiwan, [4] Inst. of Space. Sci., NCU in Taiwan, [5] Tokyo Metro. College of Aero. Eng.

Study of seismo-electromagnetics has been developed for a few decades. In particular, ionospheric disturbances associated with earthquakes have been proposed since the 1980s and a few of recent publications showed strong evidence for the existence of seismo-ionospheric disturbances. Consequently, we tried to investigate whether atmospheric disturbances associated with earthquakes also exist, by measuring anomalous transmission of VHF electromagnetic waves. The VHF electromagnetic waves scattered by the atmospheric or ionospheric disturbances beyond the line-of-sight distance could be observed through our one-year observation. For our analysis, we used one data a day that was derived by the variation of the amplitude of the received VHF electromagnetic waves during the midnight after a wavelet noise reduction was performed. When we compare the variation of such a one-day data with the upper bound derived by the previous 14-days running median of the one-day data and the associated 2 times inter-quartile range, some signatures over the upper bound might appear a few days before earthquakes (more than M4.8) that occur between transmitter and receiver. We finally obtained 11 signals that appeared a few days before 15 earthquakes. Furthermore, we analyzed the polarization of the 11 signals, carefully comparing the anomalous transmission caused by foEs, meteor shower and so on. It is concluded that there also was a possibility of the atmospheric disturbance associated with the earthquakes such as the ionospheric one because the transmission path was not the ionosphere but the atmosphere due to no change of the polarization of the received VHF electromagnetic waves.

Acknowledgments: The authors would like to thank Mr. M. Ikeda and Mr. S. Muramatsu for their help, useful comments and advice.