

Redetermination of hypocenters in SW Japan using the data of JMA and universities in 1975 and 1976

Yuzo Ishikawa[1], Kiyoshi Ito[2], Hiroshi Katao[3], Tadashi Konomi[4], Kunihiko Watanabe[3], Shozo Kimura[5], Masao Nakamura[6], Katsumi Miura[7], Tooru Ooida[8]

[1] Meteorological Res. Inst., [2] Disas. Prev. Res. Inst, Kyoto Univ., [3] RCEP, DPRI, Kyoto Univ., [4] Research Center for Earthquake Prediction, DPRI, Kyoto Univ., [5] Earthq.Obs.,Sci.,Kochi Univ., [6] Wakayama Obs.,Observation Center,ERI,Univ. of Tokyo., [7] ERI,Univ. of Tokyo, [8] RCSV, Nagoya Univ.

Arrival time data of National universities were combined with that of JMA data for getting more detail seismic activity data. The data of Tottori, Abuyama, and Tokushima seismic observatories of Kyoto university, Wakayama seismic observatory of ERI, Tochi seismic observatory of Kochi university, and Takayama seismic observatory of Nagoya university for 1975 and 1976 were gathered and inputed. The number of hypocenters determined were 20 times as that of JMA catalogue.