

# Geophysical meanings and geodetic aspects of the generalized Bouguer anomaly

# Kyozo Nozaki[1]

[1] OYO

The author has proposed a new concept of the generalized Bouguer anomaly, which is defined upon the datum level of an arbitrary equipotential surface (e.g. Nozaki, K., 2003). In this manuscript, the author presents the geophysical meanings and geodetic aspects of the generalized Bouguer anomaly. The main viewpoints are: (1) How should be the reference gravity/density field of the geophysical gravity anomaly?, (2) What is the relation between the equation of the generalized Bouguer anomaly and the fundamental equation of physical geodesy?, etc.

The main purpose of introducing the idea of the generalized Bouguer anomaly is to supplement the defects of the classical Bouguer anomaly (e.g. Heiland, 1946), in which the comparison between the observed and the reference gravities is done on a common datum level of the geoid; while the normal gravity (Heiskanen and Moritz, 1967) has been adopted as the reference gravity in defining the Bouguer anomaly today, in which the comparison is done at two distinct levels: the geoid and the normal ellipsoid. Hence, the geophysical meaning of the Bouguer anomaly today, which should be originally aimed at investigating subsurface density anomaly, is in confusion.

In defining a so-called 'anomaly', one can choose in a sense any one of references according to the purposes. From a view point of geophysical gravity anomaly that is aimed at dealing with the density anomaly, it is natural to select a reference density field with a constant/mean density, separating the actual earth's density field into [the mean] and [the deviatoric]. Accordingly, the reference gravity field of the generalized Bouguer anomaly is to be the Prey-reduced normal one within the mean density field, in which the values on the normal ellipsoid surface coincide with those of the normal gravity field. By adopting the Prey-reduced normal gravity field, one can define a specific datum level of gravity reduction (i.e. density-free datum level), and can realize the density-free Bouguer anomaly upon the density-free datum level, that is, the generalized Bouguer anomaly becomes free from the Bouguer reduction density at the density-free specific datum level. The author demonstrates, in the presentation, this fact leads to the results that (1) the idea of the generalized Bouguer anomaly removes the ambiguities in the definition of the Bouguer anomaly and (2) the generalized Bouguer anomaly at the density-free datum level is consistent with the geodetic frame, which totally mediate between the geophysical and geodetic gravity anomalies.

Acknowledgements: The author expresses his sincere thanks to Profs. Emeritus Shozaburo Nagumo, Yukio Hagiwara of the University of Tokyo, Profs. Shuhei Okubo, Yoshio Fukao of the University of Tokyo, and Drs. Masaru Kaidzu, Yuki Kuroishi of the Geographical Survey Institute for their helpful comments.

## References:

- [1] Nozaki (2003): The generalized Bouguer anomaly, submitted to J. Geod. (Oct. 2003).
- [2] Moritz (1990): The figure of the earth, WICHMANN, pp.279.
- [3] Heiskanen and Moritz (1967): Physical geodesy, WH Freeman & Company, pp.364.
- [4] Hackney and Featherstone (2003): Geodetic versus geophysical perspectives of the 'gravity anomaly', Geophys. J. Int., 154,35-43.
- [5] Vanicek et al. (2003): New views of the spherical Bouguer gravity anomaly, submitted to Geophys. J. Int., (personal comm.).
- [6] IUGG/IAG (1967): The geodetic reference system 1967.
- [7] Heiland (1946): Geophysical exploration Prentice-Hall, Inc., pp.1013.
- [8] Kuhn (2003): Geoid determination with density hypotheses from isostatic models and geological information, J. Geod., 77, 50-65.