

## GPS Monitoring around Tokai Region

# Sayaka Uchiumi[1]; Yasuko Shimizu[1]; Yasushi Harada[2]; Mikio Satomura[3]; Teruyuki Kato[4]; Atsumi Kumamoto[5]; Seiichi Shimada[6]; Jun'ichi Fukuda[7]; Shin'ichi Miyazaki[8]; Makiko Iwakuni[4]; Fumiaki Kimata[9]; Takeshi Matsushima[10]; Toshiyasu Nagao[11]; Keizo Sayanagi[12]; Shigekazu Kusumoto[13]

[1] Shizuoka Univ.; [2] School of Marine Sci. and Tech., Tokai Univ.; [3] Fac. of Science, Shizuoka Univ.; [4] Earthq. Res. Inst., Univ. Tokyo; [5] Biology and Geosciences Sci., Shizuoka Univ.; [6] NIED; [7] ERI, Univ. of Tokyo; [8] ERI; [9] Res. Center Seis. & Volcanology, Graduate school of Environ., Nagoya Univ.; [10] SEVO, Kyushu Univ.; [11] Earthquake Prediction Res. Center, Tokai Univ.; [12] IORD, Tokai Univ.; [13] School of Marine Sci. & Tech., Tokai Univ.

New dense GPS network around Tokai region is established by Japanese university consortium for GPS Research (JUNCO) which mainly consists of University of Tokyo (Earthquake Research Institute), Shizuoka University, University of Tokai and Kyushu University. The region is widely noticed for slow slip (or slow earthquake) which is occurring continuously more than five years from 2000 in the nine years history of GPS network(GEONET) by Geographical Survey Institute (GSI). We have installed about 60 more GPS stations in the area where about 40 GPS stations by GSI are available. Using the combined 100 GPS stations, more detailed feature of the slow slip in Tokai region is expected to observe. We will show some preliminary results of the combined new GPS network which is now 2.5 times denser than previous one.