

Comprehensive research of the Chuseki-so in the Tokyo and Nakagawa Lowlands, Kanto Plain, at the fourth year stage

Katsumi Kimura[1]; Susumu Tanabe[2]; Toshimichi Nakanishi[2]; Yoshiro Ishihara[3]; Chikako Eto[4]; Mieko Uchiyama[5]; Tomio Inazaki[2]; Takato Takemura[6]; Takahiro Matsumoto[7]; Masanobu Oda[8]; Rei Nakashima[9]; Shoichi Hachinohe[10]; Toshio Nakayama[11]

[1] GSJ,AIST; [2] GSJ, AIST; [3] Fukuoka Univ.; [4] Earth System Sci., Fukuoka Univ.; [5] GSJ/AIST; [6] AIST; [7] Soil, Civil&Enviroment, Saitama University; [8] Civil and Environmental Engineering, Saitama Univ.; [9] Institute of Geoscience, AIST; [10] Center for Envir. Sci., Saitama; [11] Institute of Civil Engineering of T.M.G.

The Tokyo and Nakagawa Lowlands are underlain by the Latest Pleistocene to Holocene strata, called as Chuseki-so deposits, filling a buried incised valley formed during the last glacial age. The science project has been doing as follows; 1) establishing the standard of stratigraphy and physical property of the Chuseki-so deposits, 2) detailed structural image of the ground, such as the shape of the buried incised valley and the internal structure of the Chuseki-so deposits by the S wave reflection study, 3) collecting about 6000 numbers of existing borehole log data surveyed for building construction and establishing a 3-D geological structural model.

We will introduce the recent study results at the fourth year and the outline of a next medium-range plan.