

## Direct comparison of MSTIDs observed at different latitudes

# Tetsuro Ishida[1]; Keisuke Hosokawa[1]; Satoshi Shimizu[1]; Natsuo Sato[2]; Akira Sessai Yukimatu[3]; Ichiro Tomizawa[4]; Takashi Shibata[1]; Akinori Saito[5]; Nozomu Nishitani[6]

[1] Univ. of Electro-Communications; [2] NIPR; [3] NIPR (SOKENDAI, Polar Science); [4] Sugadaira Space Radio Obs., Univ. of Electro-Comm.; [5] Dept. of Geophysics, Kyoto Univ.; [6] STELAB, Nagoya Univ.

It is commonly believed that Medium-Scale Traveling Ionospheric Disturbance (MSTIDs) are generated in the auroral region and then propagate radially toward the mid-latitudes. However, one to one relationship between the MSTIDs observed at the auroral and mid-latitudes has not been confirmed by substantial measurements. In order to clarify whether or not sources of the MSTIDs observed at the auroral and mid-latitudes are common, we directly compared periods of the MSTIDs observed almost

simultaneously at two different latitudes. SuperDARN radars in Alaska were employed for detecting MSTIDs at the auroral latitudes, and HF Doppler sounding system operated by UEC and SuperDARN radar in Hokkaido were used at the mid-latitudes.

We have detected a few events of equatorward propagating MSTIDs observed simultaneously in Alaska and Japan. Then, it was found that dominant period of the MSTIDs observed at auroral and mid-latitude are very similar, suggesting the long-distance trip of

the MSTIDs from the auroral to mid-latitudes.