

## Atmospheric escape from the early Martian atmosphere

# Naoki Terada[1]; Y. Kulikov[2]; H. Lammer[3]; M. Khodachenko[3]; H. Lichtenegger[3]

[1] NICT/JST; [2] PGI, Russian Academy of Sciences; [3] SRI, Austrian Academy of Sciences

The evolution of the Martian atmosphere inventory is influenced by thermal and non-thermal atmospheric loss into space. In order to study how much of the Martian atmosphere could have been lost to space over the planet's history and discuss the effect of the total atmospheric loss on the evolution of the climate and hydrosphere of Mars, we apply global 3-D magnetohydrodynamic (MHD) and particle simulation models of the solar wind interaction with the upper atmosphere of Mars. For reconstructing the past solar radiation and solar wind environment we use data from the observation of solar proxies with different ages. For modeling thermosphere over the Martian history we apply a diffusive-gravitational equilibrium and thermal balance model which investigate the heating of the thermosphere by photodissociation and ionization processes, due to exothermic chemical reactions and cooling by CO<sub>2</sub> IR emission in the 15 micrometer band. We present simulation results of the solar wind interaction with the XUV-heated, highly expanded upper atmosphere of early Mars as well as estimates of non-thermal ion escape rate, and compare them with those for the present Mars.