

Spatial distribution of ionospheric disturbances prior to the Sumatra earthquakes

Masahide Nishihashi[1]; Katsumi Hattori[2]; Jann-Yenq Liu[3]; Sarmoko Saroso[4]; Djedi S. Widarto[5]

[1] Geosys. and Biosys. Sci. Div., Graduate School of Sci. and Tech., Chiba Univ.; [2] Chiba University; [3] Institute of Space Science, National Central University, Taiwan ; [4] LAPAN, Indonesia; [5] LIPI, Indonesia

<http://www-es.s.chiba-u.ac.jp/geoph/ulf/>

Many anomalous electromagnetic phenomena possibly associated with large earthquakes have been reported. Recent researchers found apparent reductions in the ionospheric total electron content (TEC) within 1 - 5 days prior to M 5.0- earthquakes in Taiwan, as exemplified by the 1999 Chi-Chi earthquake (Mw7.6) and Chia-Yi earthquake (ML6.4). However, these studies did not consider the simultaneous data sets from other sites outside of Taiwan. In this paper, we investigate whether the anomalies observed during these earthquakes were local or global phenomena using TEC data observed by ground-based GPS receivers and NmF2 data obtained from ionosondes in Taiwan and Japan. We also used the global ionosphere maps (GIM) published by the Center for Orbit Determination in Europe (CODE) in order to detect anomalies in various locations throughout the globe. In this result, we have found that the anomaly four days before the Chi-Chi earthquake was global phenomenon induced by the geomagnetic disturbance. However, we have recognized three anomalies to be local phenomena. They occurred (1) three days before the Chi-Chi earthquake, (2) one day before the Chia-Yi earthquake, and (3) three days before the Chia-Yi earthquake. For all of these three anomalies, the ionospheric-disturbed areas were localized around Taiwan, and did not spread all the way to Tokyo. The disturbed areas were within a 2200 km radius and seem to be much smaller. We have applied the proposed algorithm to the 2004 - 2007 Sumatra earthquakes. The detail results will be shown in the presentation.