P221-034 Room: 302 Time: May 26 12:11-12:12

## Craters Recognition Tool on Digital Terrain Model from LISM/TC, KAGUYA

# Noriaki Asada[1]; Naoto Harada[1]; Naru Hirata[1]; Hirohide Demura[1]; Jun'ichi Haruyama[2]; Makiko Ohtake[2]; Tsuneo Matsunaga[3]; Yasuhiro Yokota[2]; Tomokatsu Morota[2]; Chikatoshi Honda[4]; Yoshiko Ogawa[3]; Haruyama Jun-ichi LISM Working Group[5]

[1] Univ. of Aizu; [2] ISAS/JAXA; [3] NIES; [4] ISAS; [5] -

http://www.u-aizu.ac.jp

A tool for automatic recognition of craters and linear features on Digital Terrain Model is shown. This core technique is an application of generalized Hough transform, where Ellipsoid-modeled Hough Transform (EHT) is adopted for circular features[1]. Pre-processing and post-processing (refitting) has been also optimized and a function of extracting linear features such as wrinkle ridges, grabens, and rills has been added. This tool is prepared for LISM [2][3][4] / TC (Terrain Camera [5]), Kaguya (SELENE) [6]. Verification of this tool with MOLA DTM and preliminary results for Kaguya DTM are shown. EHT has shown a strong tolerance for noises and modifications; overlaid by other craters, partly eroded from them, large collapse of crater rims.

Craters (circular features) recognition is a hot field for planetary science and computer science. Various approaches have been proposed, and they can be classified by input data; images and DTMs. The former researches are Hough Transform [7], Fuzzy Hough Transform [8], Template matching [9], or Learning algorithm [10]. On the other hand, a few researches have been found for the latter one, only that of Hough Transform [11][12]. There typical crater recognition rates range from 60 percents to 80 percents. Our tool shows the similar value and the verification with the size-frequency distribution of craters from a viewpoint of handling craters in planetary science.

Comparison of size-frequency distribution of craters between automatic results and manual one shows that the shape of plots is reliable in spite of a little systematic decreasing.

## References:

- [1] Matsumoto, N. et al. (2005) LPSC XXXVI, #1995.
- [2] Haruyama, J. et al. (2008) Earth Planets and Space, in press.
- [3] Ohtake, M. et al.(2008) Advances in Space Research, in press.
- [4] Matsunaga, T. et al. (2001) Proc. SPIE, 4151, 32-39.
- [5] Haruyama, J. et al. (2006) LPSC XXXVII, #1132.
- [6] Kato, M. et al. (2007) AGU Fall Meeting 2007, #P44A-01.
- [7] Michael, G. G. et al. (2003), PSS 51, 563.
- [8] Sawabe, Y. et al. (2006), Advances in Space Research 37, 21.
- [9] Bandeira, L. P. C. et al. (2006) 1st Intl. Conf. on Impact Cratering in the Solar System, 17.
- [10] Honda, R. et al. (2002) Progress of Discovery Science, Springer Verlag, 395.
- [11] Bruzzone, L. et al. (2004), Proc. ESA-EUSC 2004, 13.
- [12] Alves, E. I. (2003), EGS-AGU-EUG, #8974