

## Observation of anomalous ion acceleration around the moon by MAP-PACE

# Shoichiro Yokota[1]; Yoshifumi Saito[2]; Kazushi Asamura[1]; Takaaki Tanaka[3]; Masaki Nishino[4]; Tadateru Yamamoto[4]; Hideo Tsunakawa[5]; Toshio Terasawa[6]; Yoshifumi Saito KAGUYA MAP-PACE Team[7]; TSUNAKAWA, Hideo KAGUYA MAP-LMAG Team[7]

[1] ISAS/JAXA; [2] ISAS; [3] Dept. of Earth and Planetary Sci., Tokyo Univ.; [4] Earth and Planetary Sci., Univ. of Tokyo; [5] Dept. Earth Planet. Sci., Tokyo TECH; [6] Dept. Phys., Tokyo Tech.; [7] -

We report the ion acceleration around the moon observed by the MAP instrument onboard KAGUYA (SELENE). KAGUYA is a Japanese lunar orbiter that was launched on 14 September 2007 and now is at  $\sim 100$  km altitude. MAP consists of LMAG for the magnetic field observation and PACE for the plasma. PACE can measure the full three-dimensional distribution function of low energy electrons below 15 keV and ions below 28 keV. We investigated MAP data to find clear energy dispersion of ions near the terminator of the moon. The maximum energy of the ions is much higher than the kinetic energy of the solar wind ions, which means occurrence of remarkable ion acceleration around the moon. Primary candidates for the acceleration mechanism are (1) electric field induced by the solar wind and (2) electric field near the terminator of the lunar surface. We will also discuss possible sources of the observed high-energy ions.