

## Global water group neutral cloud model in Saturn's inner magnetosphere

# Hiroyasu Tadokoro[1]; Hiroaki Misawa[2]; Fuminori Tsuchiya[1]; Yuto Katoh[3]; Akira Morioka[4]; Mizuki Yoneda[1]

[1] Planet. Plasma Atmos. Res. Cent., Tohoku Univ.; [2] Planet. Plasma Atmos. Res. Cent., Tohoku Univ.  
; [3] Grad. Sch. Sci, Tohoku Univ.; [4] Planet. Plasma Atmos. Res. Cent., Tohoku Univ.

Water group neutral particles are considered to dominate the dynamics of Saturn's inner magnetosphere since neutral density in Saturn's inner magnetosphere is approximately ten times greater than plasma density. Therefore, it is important to understand a neutral distribution in Saturn's inner magnetosphere. Cassini observations have revealed that icy moon Enceladus ( $L \sim 3.94$ ) is highly active with a plume of water from its south polar region (Porco et al., 2006).

Solving a kinetic equation by referring to Ip (1997), we have derived a global distribution of water group neutral cloud in Saturn's inner magnetosphere using a Monte-Carlo procedure. We also used plasma parameters such as ion densities and electron temperatures, which depend on chemical reactions, based on Cassini observations. We consider sputtering from Enceladus, Tethys, Dione, and Rhea, and the plume from the south pole of Enceladus as release processes of water molecules. In this presentation, we will show calculational results and discuss global neutral distributions, especially longitudinal dependence due to hot electrons.