

AAS001-03

会場: 201B

時間: 5月28日09:30-09:45

上部対流圏におけるCO₂の南北勾配の長期変化

A long-term change in north-to-south gradient of CO₂ in the upper troposphere

松枝 秀和^{1*}, 町田 敏暢², 澤 庸介¹

Hidekazu Matsueda^{1*}, Toshinobu Machida², Yousuke Sawa¹

¹気象研究所 地球化学研究部, ²国立環境研究所

¹Meteorological Research Institute, ²NIES

Since April 1993, flask samplings for the JAL project had been made biweekly using a Boeing 747 passenger aircraft over the western Pacific between Japan and Australia. The air samples collected at 8-13 km were precisely measured for concentrations of CO₂ and other trace gases. This JAL data is successful for completely describing time-series variations of upper tropospheric CO₂ for 12 latitudinal bands between 30N and 30S during the past 16 years.

Using the JAL observational record, we could analyze several characteristics of long-term changes in the spatial CO₂ distributions in the upper troposphere. In the present study, we present the latitudinal distribution of CO₂, focusing on a gradual change of north-to-south gradient in the upper troposphere. This change is also evidenced when the CO₂ data observed in 1984 and 1985 by Nakazawa et al. (1991) are compared. It indicates a large impact of the increasing fossil fuel emissions on the global-scale CO₂ distribution.

Based on the extrapolated relation with the anthropogenic emissions, we deduced the natural latitudinal distribution of CO₂ in the upper troposphere when the fossil fuel emission is near zero during the pre-industrial age. Two key features of the natural distribution include: (1) CO₂ concentrations in the Northern Hemisphere that are lower than those in the Southern Hemisphere; (2) CO₂ concentration differences that are higher in the tropics than those currently measured.

キーワード: 二酸化炭素, 上部対流圏, 南北勾配

Keywords: CO₂, upper troposphere, north-to-south gradient