

AAS003-01

会場: 301B

時間: 5月27日10:45-11:00

超伝導サブミリ波リム放射サウンダ(SMILES)の概要と初期成果

Overview and Early Results of the Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES)

塩谷 雅人^{1*}, 高柳 昌弘², 村山 泰啓³

Masato Shiotani^{1*}, Masahiro Takayanagi², Yasuhiro Murayama³

¹京都大学, ²宇宙航空研究開発機構, ³情報通信研究機構

¹Kyoto University, ²JAXA, ³NICT

The Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES) was developed to be aboard the Japanese Experiment Module (JEM) on the International Space Station (ISS) under the cooperation of the Japan Aerospace Exploration Agency (JAXA) and the National Institute of Information and Communication Technology (NICT). SMILES was successfully launched by the H-II/B rocket with the H-II Transfer Vehicle (HTV) on September 11, 2009, was attached to JEM on September 25, and started atmospheric observations on October 12 (All dates in JST). Mission objectives are: i) Space demonstration of 4-K mechanical cooler and super-conductive mixer for the submillimeter limb-emission sounding in the frequency bands of 624.32- 627.32 GHz and 649.12- 650.32 GHz, and ii) global measurements with its high sensitivity for atmospheric minor constituents in the middle atmosphere (O₃, HCl, ClO, HO₂, HOCl, BrO, O₃ isotopes, HNO₃, CH₃ CN, etc), contributing to the atmospheric sciences.

Now SMILES is in the operational mode and has been performing global observations about 100 points per each track of ISS, except for any restrictions due to the ISS operation. After the data processing, global distributions of about ten kinds of atmospheric minor constituents related to the ozone chemistry have been revealed, which will contribute to various issues of atmospheric science. Since ISS has a circular orbit with an inclination angle of 51.6 degrees, no observation in the polar latitude is available. To measure northern high-latitude regions the antenna beam is tilted 45 degrees left from the direction of orbital motion, enabling SMILES to observe latitudes from 38°S to 65°N. Another important aspect of the SMILES observation is that the ISS orbit is not sun synchronous. Therefore, SMILES can measure distributions of the above mentioned minor species at different local time, and consequently SMILES can provide information on the diurnal cycle of them.

The level 2 processing has been run to retrieve vertical profiles of these species. At the same time we have been conducting preliminary data quality check by comparing the SMILES data with other existing data source. Results from SMILES have demonstrated its high potential to observe the atmospheric minor constituents in the stratosphere and the mesosphere. We hope that these outstanding data from SMILES can make a breakthrough in the middle atmosphere science.

キーワード:衛星観測,国際宇宙ステーション,成層圏オゾン,中層大気

Keywords: satellite observations, International Space Station, stratospheric ozone, middle atmosphere