

AEM011-P07

会場:コンベンションホール

時間: 5月26日17:15-18:45

ノルウェー, トロムソにおけるナトリウムライダー観測

Measurements by a sodium lidar at Tromsø, Norway

津田 卓雄^{1*}, 野澤 悟徳¹, 川原 琢也², 川端 哲也¹, 大山 伸一郎¹, 藤井 良一¹, 小川 泰信³,
斎藤 徳人⁴, 和田 智之⁴, Brekke Asgeir⁵, Hall Chris M.⁵

Takuo Tsuda^{1*}, Satonori Nozawa¹, Takuya Kawahara², Tetsuya Kawabata¹,
Shin-ichiro Oyama¹, Ryoichi Fujii¹, Yasunobu Ogawa³, Norihito Saito⁴, Satoshi Wada⁴,
Brekke Asgeir⁵, Hall Chris M.⁵

¹名古屋大学太陽地球環境研究所, ²信州大学工学部, ³国立極地研究所, ⁴理化学研究所, ⁵トロムソ大学

¹STEL, Nagoya Univ., ²Faculty of Engineering, Shinshu Univ., ³National Institute of Polar Research, ⁴RIKEN,
⁵Univ. of Tromsø

Of vital importance is to qualify significance of the magnetospheric energy input to the upper atmosphere in the polar region, in order to obtain better understanding of the energy process from the Sun to the Earth. Many observations by Incoherent Scatter (IS) radars and Fabry-Perot Interferometers (FPIs) demonstrated neutral wind variations in the polar lower thermosphere during geomagnetically active intervals. These results suggest that the neutral wind dynamics is influenced by the thermal (due to Joule and particle heating) and/or kinetic (due to ion drag) energy input from the magnetosphere. A case study, which is based on IS radar data, determined contributions of the Joule heating and the ion drag on an observed neutral wind variation, and intended that the Joule heating was a major important factor. However, there was no observational evidence of the expected temperature increase due to the Joule heating. To detect the temperature increase, it is worthwhile to conduct simultaneous observations by a sodium temperature lidar as well as an IS radar.

Since 2007, we have developed a new sodium lidar and prepared to install this lidar at Tromsø, European Incoherent SCATter (EISCAT) radar site (69.6 deg N, 19.2 deg E). The lidar system has now been nearly completed. Using the lidar, we performed test observations in November and December 2009 at Wako, RIKEN (35.8 deg N, 139.6 deg E), and detected backscatter signals from the sodium layers (80-100 km in height). The lidar will be ready for operation at Tromsø in February 2010. In this presentation, we will briefly introduce the new sodium lidar system, and then present results of the test observations at Wako. Furthermore, we will report first measurements by the lidar in February-March 2010, and discuss accuracy of the obtained temperature data.

キーワード:ナトリウムライダー, トロムソ, EISCAT

Keywords: Sodium lidar, Tromsø, EISCAT